



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁵ : A61K 48/00 | A1 | (11) International Publication Number: WO 94/05333 (43) International Publication Date: 17 March 1994 (17.03.94) |
| (21) International Application Number: PCT/US93/08101 (22) International Filing Date: 27 August 1993 (27.08.93) (30) Priority data: 07/939,855 2 September 1992 (02.09.92) US 08/007,997 21 January 1993 (21.01.93) US 08/063,167 17 May 1993 (17.05.93) US (71) Applicant: ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US). (72) Inventors: BENNET, C., Frank ; 2613 Valewood, Carls- bad, CA 92008 (US). MIRABELLI, Christopher, K. ; 6 Pine Street, Dover, MA 02030 (US). (74) Agent: LICATA, Jane, Massey; Woodland Falls Corporate Park, 210 Lake Drive East, Suite 201, Cherry Hill, NJ 08002 (US). | | (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: OLIGONUCLEOTIDE MODULATION OF CELL ADHESION | | |
| (57) Abstract Compositions and methods are provided for the treatment and diagnosis of diseases amenable to treatment through modulation of the synthesis or metabolism of intercellular adhesion molecules. In accordance with preferred embodiments, oligonucleotides are provided which are specifically hybridizable with nucleic acids encoding intercellular adhesion molecule-1, vascular cell adhesion molecule-1, and endothelial leukocyte adhesion molecule-1. The oligonucleotide comprises nucleotide units sufficient in identity and number to effect said specific hybridization. In other preferred embodiments, the oligonucleotides are specifically hybridizable with a transcription initiation site, a translation initiation site, 5'-untranslated sequences, 3'-untranslated sequences, and intervening sequences. Methods of treating animals suffering from disease amenable to therapeutic intervention by modulating cell adhesion proteins with an oligonucleotide specifically hybridizable with RNA or DNA corresponding to one of the foregoing proteins are disclosed. Methods for treatment of diseases responding to modulation cell adhesion molecules are disclosed. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|---------------------------------------|----|--------------------------|
| AT | Austria | FR | France | MR | Mauritania |
| AU | Australia | GA | Gabon | MW | Malawi |
| BB | Barbados | GB | United Kingdom | NE | Niger |
| BE | Belgium | GN | Guinea | NL | Netherlands |
| BF | Burkina Faso | GR | Greece | NO | Norway |
| BG | Bulgaria | HU | Hungary | NZ | New Zealand |
| BJ | Benin | IE | Ireland | PL | Poland |
| BR | Brazil | IT | Italy | PT | Portugal |
| BY | Belarus | JP | Japan | RO | Romania |
| CA | Canada | KP | Democratic People's Republic of Korea | RU | Russian Federation |
| CF | Central African Republic | KR | Republic of Korea | SD | Sudan |
| CG | Congo | KZ | Kazakhstan | SE | Sweden |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovak Republic |
| CM | Cameroon | LU | Luxembourg | SN | Senegal |
| CN | China | LV | Latvia | TD | Chad |
| CS | Czechoslovakia | MC | Monaco | TC | Togo |
| CZ | Czech Republic | MG | Madagascar | UA | Ukraine |
| DE | Germany | ML | Mali | US | United States of America |
| DK | Denmark | MN | Mongolia | UZ | Uzbekistan |
| ES | Spain | | | VN | Viet Nam |
| FI | Finland | | | | |

OLIGONUCLEOTIDE MODULATION OF CELL ADHESION

FIELD OF THE INVENTION

This invention relates to diagnostics, research reagents and therapies for disease states which respond to modulation of the synthesis or metabolism of cell adhesion molecules. In particular, this invention relates to antisense oligonucleotide interactions with certain messenger ribonucleic acids (mRNAs) or DNAs involved in the synthesis of proteins that regulate adhesion of white blood cells to other white blood cells and to other cell types. Antisense oligonucleotides designed to hybridize to the mRNA encoding intercellular adhesion molecule-1 (ICAM-1), endothelial leukocyte adhesion molecule-1 (ELAM-1, also known as E-selectin), and vascular cell adhesion molecule-1 (VCAM-1) are provided. These oligonucleotides have been found to lead to the modulation of the activity of the RNA or DNA, and thus to the modulation of the synthesis and metabolism of specific cell adhesion molecules. Palliation and therapeutic effect result.

BACKGROUND OF THE INVENTION

Inflammation is a localized protective response elicited by tissues in response to injury, infection, or tissue destruction resulting in the destruction of the infectious or injurious agent and isolation of the injured tissue. A typical inflammatory response proceeds as follows: recognition of an antigen as foreign or recognition of tissue damage, synthesis and release of soluble inflammatory mediators, recruitment of inflammatory cells to the site of infection or tissue damage, destruction and removal of the invading organism or damaged tissue, and deactivation of the system once the invading organism or damage has been resolved. In many human diseases with an inflammatory component, the normal, homeostatic mechanisms which attenuate the inflammatory responses are defective, resulting in damage and destruction of normal tissue.

Cell-cell interactions are involved in the activation of the immune response at each of the stages described above. One of the earliest detectable events in a normal inflammatory

- 2 -

response is adhesion of leukocytes to the vascular endothelium, followed by migration of leukocytes out of the vasculature to the site of infection or injury. The adhesion of these leukocytes, or white blood cells, to vascular endothelium is an obligate step in the migration out of the vasculature. Harlan, J.M., *Blood* 1985, 65, 513-525. In general, the first inflammatory cells to appear at the site of inflammation are neutrophils followed by monocytes, and lymphocytes. Cell-cell interactions are also critical for propagation of both B-lymphocytes and T-lymphocytes resulting in enhanced humoral and cellular immune responses, respectively.

The adhesion of white blood cells to vascular endothelium and other cell types is mediated by interactions between specific proteins, termed "adhesion molecules," located on the plasma membrane of both white blood cells and vascular endothelium. The interaction between adhesion molecules is similar to classical receptor ligand interactions with the exception that the ligand is fixed to the surface of a cell instead of being soluble. The identification of patients with a genetic defect in leukocyte adhesion has enabled investigators to identify a family of proteins responsible for adherence of white blood cells. Leukocyte adhesion deficiency (LAD) is a rare autosomal trait characterized by recurrent bacterial infections and impaired pus formation and wound healing. The defect was shown to occur in the common B-subunit of three heterodimeric glycoproteins, Mac-1, LFA-1, and p150,95, normally expressed on the outer cell membrane of white blood cells. Anderson and Springer, *Ann. Rev. Med.* 1987, 38, 175-194. Patients suffering from LAD exhibit a defect in a wide spectrum of adherence-dependent functions of granulocytes, monocytes, and lymphocytes. Three ligands for LFA-1 have been identified, intercellular adhesion molecules 1, 2 and 3 (ICAM-1, ICAM-2 and ICAM-3). Both Mac-1 and p150,95 bind complement fragment C3bi and perhaps other unidentified ligands. Mac-1 also binds ICAM-1.

Other adhesion molecules have been identified which are involved in the adherence of white blood cells to vascular

- 3 -

endothelium and subsequent migration out of the vasculature. These include endothelial leukocyte adhesion molecule-1 (ELAM-1), vascular cell adhesion molecule-1 (VCAM-1) and granule membrane protein-140 (GMP-140) and their respective receptors.

5 The adherence of white blood cells to vascular endothelium appears to be mediated in part if not *in toto* by the five cell adhesion molecules ICAM-1, ICAM-2, ELAM-1, VCAM-1 and GMP-140. Dustin and Springer, *J. Cell Biol.* 1987, 107, 321-331. Expression on the cell surface of ICAM-1, ELAM-1, VCAM-1 and
10 GMP-140 adhesion molecules is induced by inflammatory stimuli. In contrast, expression of ICAM-2 appears to be constitutive and not sensitive to induction by cytokines. In general, GMP-140 is induced by autocooids such as histamine, leukotriene B₄, platelet activating factor, and thrombin. Maximal expression
15 on endothelial cells is observed 30 minutes to 1 hour after stimulation, and returns to baseline within 2 to 3 hours. The expression of ELAM-1 and VCAM-1 on endothelial cells is induced by cytokines such as interleukin-1 β and tumor necrosis factor, but not gamma-interferon. Maximal expression of ELAM-1 on the
20 surface of endothelial cells occurs 4 to 6 hours after stimulation, and returns to baseline by 16 hours. ELAM-1 expression is dependent on new mRNA and protein synthesis. Elevated VCAM-1 expression is detectable 2 hours following treatment with tumor necrosis factor, is maximal 8 hours
25 following stimulation, and remains elevated for at least 48 hours following stimulation. Rice and Bevilacqua, *Science* 1989, 246, 1303-1306. ICAM-1 expression on endothelial cells is induced by cytokines interleukin-1 tumor necrosis factor and gamma-interferon. Maximal expression of ICAM-1 follows that of
30 ELAM-1 occurring 8 to 10 hours after stimulation and remains elevated for at least 48 hours.

GMP-140 and ELAM-1 are primarily involved in the adhesion of neutrophils to vascular endothelial cells. VCAM-1 primarily binds T and B lymphocytes. In addition, VCAM-1 may
35 play a role in the metastasis of melanoma, and possibly other cancers. ICAM-1 plays a role in adhesion of neutrophils to vascular endothelium, as well as adhesion of monocytes and

lymphocytes to vascular endothelium, tissue fibroblasts and epidermal keratinocytes. ICAM-1 also plays a role in T-cell recognition of antigen presenting cell, lysis of target cells by natural killer cells, lymphocyte activation and proliferation, and maturation of T cells in the thymus. In addition, recent data have demonstrated that ICAM-1 is the cellular receptor for the major serotype of rhinovirus, which account for greater than 50% of common colds. Staunton et al., *Cell* 1989, 56, 849-853; Greve et al., *Cell* 1989, 56, 839-847.

Expression of ICAM-1 has been associated with a variety of inflammatory skin disorders such as allergic contact dermatitis, fixed drug eruption, lichen planus, and psoriasis; Ho et al., *J. Am. Acad. Dermatol.* 1990, 22, 64-68; Griffiths and Nickoloff, *Am. J. Pathology* 1989, 135, 1045-1053; Lisby et al., *Br. J. Dermatol.* 1989, 120, 479-484; Shiohara et al., *Arch. Dermatol.* 1989, 125, 1371-1376. In addition, ICAM-1 expression has been detected in the synovium of patients with rheumatoid arthritis; Hale et al., *Arth. Rheum.* 1989, 32, 22-30, pancreatic B-cells in diabetes; Campbell et al., *Proc. Natl. Acad. Sci. U.S.A.* 1989, 86, 4282-4286; thyroid follicular cells in patients with Graves' disease; Weetman et al., *J. Endocrinol.* 1989, 122, 185-191; and with renal and liver allograft rejection; Faull and Russ, *Transplantation* 1989, 48, 226-230; Adams et al., *Lancet* 1989, 1122-1125.

It is has been hoped that inhibitors of ICAM-1, VCAM-1 and ELAM-1 expression would provide a novel therapeutic class of anti-inflammatory agents with activity towards a variety of inflammatory diseases or diseases with an inflammatory component such as asthma, rheumatoid arthritis, allograft rejections, inflammatory bowel disease, various dermatological conditions, and psoriasis. In addition, inhibitors of ICAM-1, VCAM-1, and ELAM-1 may also be effective in the treatment of colds due to rhinovirus infection, AIDS, Kaposi's sarcoma and some cancers and their metastasis. To date, there are no known therapeutic agents which effectively prevent the expression of the cellular adhesion molecules ELAM-1, VCAM-1 and ICAM-1. The use of neutralizing monoclonal antibodies against ICAM-1 in

- 5 -

animal models provide evidence that such inhibitors if identified would have therapeutic benefit for asthma; Wegner et al., *Science* 1990, 247, 456-459, renal allografts; Cosimi et al., *J. Immunol.* 1990, 144, 4604-4612, and cardiac allografts; 5 Isobe et al., *Science* 1992, 255, 1125-1127. The use of a soluble form of ICAM-1 molecule was also effective in preventing rhinovirus infection of cells in culture. Marlin et al., *Nature* 1990, 344, 70-72.

Current agents which affect intercellular adhesion 10 molecules include synthetic peptides, monoclonal antibodies, and soluble forms of the adhesion molecules. To date, synthetic peptides which block the interactions with VCAM-1 or ELAM-1 have not been identified. Monoclonal antibodies may prove to be useful for the treatment of acute inflammatory 15 response due to expression of ICAM-1, VCAM-1 and ELAM-1. However, with chronic treatment, the host animal develops antibodies against the monoclonal antibodies thereby limiting their usefulness. In addition, monoclonal antibodies are large proteins which may have difficulty in gaining access to the 20 inflammatory site. Soluble forms of the cell adhesion molecules suffer from many of the same limitations as monoclonal antibodies in addition to the expense of their production and their low binding affinity. Thus, there is a long felt need for molecules which effectively inhibit 25 intercellular adhesion molecules. Antisense oligonucleotides avoid many of the pitfalls of current agents used to block the effects of ICAM-1, VCAM-1 and ELAM-1.

PCT/US90/02357 (Hession et al.) discloses DNA sequences encoding Endothelial Adhesion Molecules (ELAMs), 30 including ELAM-1 and VCAM-1 and VCAM-1b. A number of uses for these DNA sequences are provided, including (1) production of monoclonal antibody preparations that are reactive for these molecules which may be used as therapeutic agents to inhibit leukocyte binding to endothelial cells; (2) production of ELAM 35 peptides to bind to the ELAM ligand on leukocytes which, in turn, may bind to ELAM on endothelial cells, inhibiting leukocyte binding to endothelial cells; (3) use of molecules

- 6 -

binding to ELAMS (such as anti-ELAM antibodies, or markers such as the ligand or fragments of it) to detect inflammation; (4) use of ELAM and ELAM ligand DNA sequences to produce nucleic acid molecules that intervene in ELAM or ELAM ligand expression
5 at the translational level using antisense nucleic acid and ribozymes to block translation of a specific mRNA either by masking mRNA with antisense nucleic acid or cleaving it with a ribozyme. It is disclosed that coding regions are the targets of choice. For VCAM-1, AUG is believed to be most likely; a
10 15-mer hybridizing to the AUG site is specifically disclosed in Example 17.

OBJECTS OF THE INVENTION

It is a principle object of the invention to provide therapies for diseases with an immunological component,
15 allografts, cancers and metastasis, inflammatory bowel disease, psoriasis and other skin diseases, colds, and AIDS through perturbation in the synthesis and expression of inflammatory cell adhesion molecules.

It is a further object of the invention to provide
20 antisense oligonucleotides which are capable of inhibiting the function of nucleic acids encoding intercellular adhesion proteins.

Yet another object is to provide means for diagnosis of dysfunctions of intercellular adhesion.

25 These and other objects of this invention will become apparent from a review of the instant specification.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 is the mRNA sequence of human intercellular adhesion molecule-1 (ICAM-1).

30 FIGURE 2 is the mRNA sequence of human endothelial leukocyte adhesion molecule-1 (ELAM-1).

FIGURE 3 is the mRNA sequence of human vascular cell adhesion molecule-1 (VCAM-1).

FIGURE 4 is a graphical representation of the
35 induction of ICAM-1 expression on the cell surface of various

- 7 -

human cell lines by interleukin-1 and tumor necrosis factor.

FIGURE 5 is a graphical representation of the effects of selected antisense oligonucleotides on ICAM-1 expression on human umbilical vein endothelial cells.

5 FIGURE 6A and 6B are a graphical representation of the effects of an antisense oligonucleotide on the expression of ICAM-1 in human umbilical vein endothelial cells stimulated with tumor necrosis factor and interleukin-1.

10 FIGURE 7 is a graphical representation of the effect of antisense oligonucleotides on ICAM-1 mediated adhesion of DMSO differentiated HL-60 cells to control and tumor necrosis factor treated human umbilical vein endothelial cells.

15 FIGURE 8 is a graphical representation of the effects of selected antisense oligonucleotides on ICAM-1 expression in A549 human lung carcinoma cells.

FIGURE 9 is a graphical representation of the effects of selected antisense oligonucleotides on ICAM-1 expression in primary human keratinocytes.

20 FIGURE 10 is a graphical representation of the relationship between oligonucleotide chain length, T_m and effect on inhibition of ICAM-1 expression.

25 FIGURE 11 is a graphical representation of the effect of selected antisense oligonucleotides on ICAM-1 mediated adhesion of DMSO differentiated HL-60 cells to control and tumor necrosis factor treated human umbilical vein endothelial cells.

30 FIGURE 12 is a graphical representation of the effects of selected antisense oligonucleotides on ELAM-1 expression on tumor necrosis factor-treated human umbilical vein endothelial cells.

FIGURE 13 is a graphical representation of the human ELAM-1 mRNA showing target sites of antisense oligonucleotides.

FIGURE 14 is a graphical representation of the human VCAM-1 mRNA showing target sites of antisense oligonucleotides.

35 FIGURE 15 is a line graph showing inhibition of ICAM-1 expression in C8161 human melanoma cells following treatment with antisense oligonucleotides complementary to ICAM-1.

- 8 -

FIGURE 16 is a bar graph showing the effect of ISIS 3082 on dextran sulfate (DSS)-induced inflammatory bowel disease.

SUMMARY OF THE INVENTION

5 In accordance with the present invention, oligonucleotides are provided which specifically hybridize with nucleic acids encoding intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1) and endothelial leukocyte adhesion molecule-1 (ELAM-1). The oligonucleotides
10 are designed to bind either directly to mRNA or to a selected DNA portion forming a triple stranded structure, thereby modulating the amount of mRNA made from the gene.

The former relationship is commonly denoted as "antisense." The oligonucleotides are able to inhibit the
15 function of RNA or DNA, either its translation into protein, its translocation into the cytoplasm, or any other activity necessary to its overall biological function. The failure of the RNA or DNA to perform all or part of its function results in failure of a portion of the genome controlling cell adhesion
20 molecules to be properly expressed, thus modulating said metabolism.

It is preferred to target specific genes for antisense attack. It has been discovered that the genes coding for ICAM-1, VCAM-1 and ELAM-1 are particularly useful for this
25 approach. Inhibition of ICAM-1, VCAM-1 and ELAM-1 expression is expected to be useful for the treatment of inflammatory diseases, diseases with an inflammatory component, allograft rejection, psoriasis and other skin diseases, inflammatory bowel disease, cancers and their metastasis, and viral
30 infections.

Methods of modulating cell adhesion comprising contacting the animal with an oligonucleotide hybridizable with nucleic acids encoding a protein capable of modulating cell adhesion are provided. Oligonucleotides hybridizable with an
35 RNA or DNA encoding ICAM-1, VCAM-1 and ELAM-1 are preferred. Methods for diagnosis are also a part of this invention.

- 9 -

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Antisense oligonucleotides hold great promise as therapeutic agents for the treatment of many human diseases. Oligonucleotides specifically bind to the complementary
5 sequence of either pre-mRNA or mature mRNA, as defined by Watson-Crick base pairing, inhibiting the flow of genetic information from DNA to protein. The properties of antisense oligonucleotides which make them specific for their target sequence also make them extraordinarily versatile. Because
10 antisense oligonucleotides are long chains of four monomeric units they may be readily synthesized for any target RNA sequence. Numerous recent studies have documented the utility of antisense oligonucleotides as biochemical tools for studying target proteins. Rothenberg et al., *J. Natl. Cancer Inst.*
15 1989, 81, 1539-1544; Zon, *G. Pharmaceutical Res.* 1988, 5, 539-549). Because of recent advances in synthesis of nuclease resistant oligonucleotides, which exhibit enhanced cell uptake, it is now possible to consider the use of antisense oligonucleotides as a novel form of therapeutics.

20 Antisense oligonucleotides offer an ideal solution to the problems encountered in prior art approaches. They can be designed to selectively inhibit a given isoenzyme, they inhibit the production of the enzyme, and they avoid non-specific mechanisms such as free radical scavenging or binding to
25 multiple receptors. A complete understanding of enzyme mechanisms or receptor-ligand interactions is not needed to design specific inhibitors.

DESCRIPTION OF TARGETS

The acute infiltration of neutrophils into the site
30 of inflammation appears to be due to increased expression of GMP-140, ELAM-1 and ICAM-1 on the surface of endothelial cells. The appearance of lymphocytes and monocytes during the later stages of an inflammatory reaction appear to be mediated by VCAM-1 and ICAM-1. ELAM-1 and GMP-140 are transiently
35 expressed on vascular endothelial cells, while VCAM-1 and ICAM-1 are chronically expressed.

- 10 -

Human ICAM-1 is encoded by a 3.3-kb mRNA resulting in the synthesis of a 55,219 dalton protein (Figure 1). ICAM-1 is heavily glycosylated through N-linked glycosylation sites. The mature protein has an apparent molecular mass of 90 kDa as determined by SDS-polyacrylamide gel electrophoresis. Staunton et al., *Cell* 1988, 52, 925-933. ICAM-1 is a member of the immunoglobulin supergene family, containing 5 immunoglobulin-like domains at the amino terminus, followed by a transmembrane domain and a cytoplasmic domain. The primary binding site for LFA-1 and rhinovirus are found in the first immunoglobulin-like domain. However, the binding sites appear to be distinct. Staunton et al., *Cell* 1990, 61, 243-354. Recent electron micrographic studies demonstrate that ICAM-1 is a bent rod 18.7 nm in length and 2 to 3 nm in diameter. Staunton et al., *Cell* 1990, 61, 243-254.

ICAM-1 exhibits a broad tissue and cell distribution, and may be found on white blood cells, endothelial cells, fibroblast, keratinocytes and other epithelial cells. The expression of ICAM-1 can be regulated on vascular endothelial cells, fibroblasts, keratinocytes, astrocytes and several cell lines by treatment with bacterial lipopolysaccharide and cytokines such as interleukin-1, tumor necrosis factor, gamma-interferon, and lymphotoxin. See, e.g., Frohman et al., *J. Neuroimmunol.* 1989, 23, 117-124. The molecular mechanism for increased expression of ICAM-1 following cytokine treatment has not been determined.

ELAM-1 is a 115-kDa membrane glycoprotein (Figure 2) which is a member of the selectin family of membrane glycoproteins. Bevilacqua et al., *Science* 1989, 243, 1160-1165. The amino terminal region of ELAM-1 contains sequences with homologies to members of lectin-like proteins, followed by a domain similar to epidermal growth factor, followed by six tandem 60-amino acid repeats similar to those found in complement receptors 1 and 2. These features are also shared by GMP-140 and MEL-14 antigen, a lymphocyte homing antigen. ELAM-1 is encoded for by a 3.9-kb mRNA. The 3'-untranslated region of ELAM-1 mRNA contains several sequence motifs ATTTA

- 11 -

which are responsible for the rapid turnover of cellular mRNA consistent with the transient nature of ELAM-1 expression.

ELAM-1 exhibits a limited cellular distribution in that it has only been identified on vascular endothelial cells. Like ICAM-1, ELAM-1 is inducible by a number of cytokines including tumor necrosis factor, interleukin-1 and lymphotoxin and bacterial lipopolysaccharide. In contrast to ICAM-1, ELAM-1 is not induced by gamma-interferon. Bevilacqua et al., *Proc. Natl. Acad. Sci. USA* 1987, 84, 9238-9242; Wellicome et al., *J. Immunol.* 1990, 144, 2558-2565. The kinetics of ELAM-1 mRNA induction and disappearance in human umbilical vein endothelial cells precedes the appearance and disappearance of ELAM-1 on the cell surface. As with ICAM-1 the molecular mechanism for ELAM-1 induction is not known.

VCAM-1 is a 110-kDa membrane glycoprotein encoded by a 3.2-kb mRNA (Figure 3). VCAM-1 appears to be encoded by a single-copy gene which can undergo alternative splicing to yield products with either six or seven immunoglobulin domains. Osborn et al., *Cell* 1989, 59, 1203-1211. The receptor for VCAM-1 is proposed to be CD29 (VLA-4) as demonstrated by the ability of monoclonal antibodies to CD29 to block adherence of Ramos cells to VCAM-1. VCAM-1 is expressed primarily on vascular endothelial cells. Like ICAM-1 and ELAM-1, expression of VCAM-1 on vascular endothelium is regulated by treatment with cytokines. Rice and Bevilacqua, *Science* 1989, 246, 1303-1306; Rice et al., *J. Exp. Med.* 1990, 171, 1369-1374. Increased expression appears to be due to induction of the mRNA.

For therapeutics, an animal suspected of having a disease which can be treated by decreasing the expression of ICAM-1, VCAM-1 and ELAM-1 is treated by administering oligonucleotides in accordance with this invention. Oligonucleotides may be formulated in a pharmaceutical composition, which may include carriers, thickeners, diluents, buffers, preservatives, surface active agents, liposomes or lipid formulations and the like in addition to the oligonucleotide. Pharmaceutical compositions may also include one or more active ingredients such as antimicrobial agents,

- 12 -

anti-inflammatory agents, anesthetics, and the like in addition to oligonucleotide.

The pharmaceutical composition may be administered in a number of ways depending on whether local or systemic treatment is desired, and on the area to be treated. Administration may be topically (including ophthalmically, vaginally, rectally, intranasally), orally, by inhalation, or parenterally, for example by intravenous drip, subcutaneous, intraperitoneal or intramuscular injection.

Formulations for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms or gloves may also be useful.

Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be desirable.

Formulations for parenteral administration may include sterile aqueous solutions which may also contain buffers, liposomes, diluents and other suitable additives.

Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, with course of treatment lasting from several days to several months or until a cure is effected or a diminution of disease state is achieved. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates.

The present invention employs oligonucleotides for use in antisense inhibition of the function of RNA and DNA corresponding to proteins capable of modulating inflammatory cell adhesion. In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid or deoxyribonucleic acid. This term includes oligomers consisting of naturally occurring bases, sugars and

- 13 -

intersugar (backbone) linkages as well as oligomers having non-naturally occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of properties such as, for example, enhanced cellular uptake and increased stability in the presence of nucleases.

Specific examples of some preferred oligonucleotides envisioned for this invention may contain phosphorothioates, phosphotriesters, methyl phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Most preferred are those with $\text{CH}_2\text{-NH-O-CH}_2$, $\text{CH}_2\text{-N(CH}_3\text{)-O-CH}_2$, $\text{CH}_2\text{-O-N(CH}_3\text{)-CH}_2$, $\text{CH}_2\text{-N(CH}_3\text{)-N(CH}_3\text{)-CH}_2$ and $\text{O-N(CH}_3\text{)-CH}_2\text{-CH}_2$ backbones (where phosphodiester is O-P-O-CH_2). Also preferred are oligonucleotides having morpholino backbone structures. Summerton, J.E. and Weller, D.D., U.S. Patent 5,034,506. In other preferred embodiments, such as the protein-nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide may be replaced with a polyamide backbone, the bases being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone. P.E. Nielsen, M. Egholm, R.H. Berg, O. Buchardt, *Science* 1991, 254, 1497. Other preferred oligonucleotides may contain alkyl and halogen-substituted sugar moieties comprising one of the following at the 2' position: OH, SH, SCH_3 , F, OCN, $\text{O(CH}_2\text{)}_n\text{NH}_2$ or $\text{O(CH}_2\text{)}_n\text{CH}_3$, where n is from 1 to about 10; C_1 to C_{10} lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl; Br; CN; CF_3 ; OCF_3 ; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH_3 ; SO_2CH_3 ; ONO_2 ; NO_2 ; N_3 ; NH_2 ; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group.

The oligonucleotides in accordance with this invention

- 14 -

preferably comprise from about 3 to about 50 nucleic acid base units. It is more preferred that such oligonucleotides comprise from about 8 to 25 nucleic acid base units, and still more preferred to have from about 12 to 22 nucleic acid base
5 units. As will be appreciated, a nucleic acid base unit is a base-sugar combination suitably bound to an adjacent nucleic acid base unit through phosphodiester or other bonds.

The oligonucleotides used in accordance with this invention may be conveniently and routinely made through the
10 well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including Applied Biosystems. Any other means for such synthesis may also be employed, however the actual synthesis of the oligonucleotides are well within the talents of the routineer. It is also well
15 known to use similar techniques to prepare other oligonucleotides such as the phosphorothioates and alkylated derivatives.

In accordance with this invention, persons of ordinary skill in the art will understand that messenger RNA identified
20 by the open reading frames (ORFs) of the DNA from which they are transcribed includes not only the information from the ORFs of the DNA, but also associated ribonucleotides which form regions known to such persons as the 5'-untranslated region, the 3'- untranslated region, and intervening sequence.
25 ribonucleotides. Thus, oligonucleotides may be formulated in accordance with this invention which are targeted wholly or in part to these associated ribonucleotides as well as to the informational ribonucleotides. In preferred embodiments, the oligonucleotide is specifically hybridizable with a
30 transcription initiation site, a translation initiation site, an intervening sequence and sequences in the 3'-untranslated region.

In accordance with this invention, the oligonucleotide is specifically hybridizable with portions of nucleic acids
35 encoding a protein involved in the adhesion of white blood cells either to other white blood cells or other cell types. In preferred embodiments, said proteins are intercellular

- 15 -

adhesion molecule-1, vascular cell adhesion molecule-1 and endothelial leukocyte adhesion molecule-1. Oligonucleotides comprising the corresponding sequence, or part thereof, are useful in the invention. For example, Figure 1 is a human intercellular adhesion molecule-1 mRNA sequence. A preferred sequence segment which may be useful in whole or in part is:

| | 5' | 3' | SEQ ID NO: |
|----|------------------------|----|------------|
| | TGGGAGCCATAGCGAGGC | | 1 |
| | GAGGAGCTCAGCGTCGACTG | | 2 |
| 10 | GACACTCAATAAATAGCTGGT | | 3 |
| | GAGGCTGAGGTGGGAGGA | | 4 |
| | CGATGGGCAGTGGGAAAG | | 5 |
| | GGGCGCGTGATCCTTATAGC | | 6 |
| | CATAGCGAGGCTGAGGTTGC | | 7 |
| 15 | CGGGGGCTGCTGGGAGCCAT | | 8 |
| | TCAGGGAGGCGTGGCTTGTG | | 13 |
| | CCTGTCCCGGGATAGGTTCA | | 14 |
| | TTGAGAAAGCTTTATTA ACT | | 16 |
| | CCCCCACC ACTTCCCCTCTC. | | 15 |

Figure 2 is a human endothelial leukocyte adhesion molecule-1 mRNA sequence. A preferred sequence segment which may be useful in whole or in part is:

| | 5' | 3' | SEQ ID NO: |
|----|--------------------------|----|------------|
| | CAATCATGACTTCAAGAGTTCT | | 28 |
| 25 | TCACTGCTGCCTCTGTCTCAGG | | 73 |
| | TGATTCTTTTGA ACTTAAAAGGA | | 74 |
| | TTAAAGGATGTAAGAAGGCT | | 75 |
| | CATAAGCACATTTATTGTC | | 76 |
| | TTTTGGGAAGCAGTTGTTCA | | 77 |
| 30 | AACTGTGAAGCAATCATGACT | | 78 |
| | CCTTGAGTGGTGCATTCAACCT | | 79 |
| | AATGCTTGCTCACACAGGCATT. | | 80 |

Figure 3 is a human vascular cell adhesion molecule-1 mRNA sequence. A preferred sequence segment which may be useful in whole or in part is:

| 5' | 3' | SEQ ID NO: |
|-----------------------|----|------------|
| CCAGGCATTTTAAGTTGCTGT | | 40 |

- 16 -

| | | |
|---|------------------------|----|
| | CCTGAAGCCAGTGAGGCCCG | 41 |
| | GATGAGAAAATAGTGAACCA | 42 |
| | CTGAGCAAGATATCTAGAT | 43 |
| | CTACACTTTTGATTTCTGT | 44 |
| 5 | TTGAACATATCAAGCATTAGCT | 45 |
| | TTTACATATGTACAAATTATGT | 46 |
| | AATTATCACTTTACTATACAAA | 47 |
| | AGGGCTGACCAAGACGGTGT. | 48 |

While the illustrated sequences are believed to be accurate, the present invention is directed to the correct sequences should errors be found. Oligonucleotides useful in the invention comprise one of these sequences, or part thereof. Thus, it is preferred to employ any of these oligonucleotides as set forth above or any of the similar oligonucleotides which persons of ordinary skill in the art can prepare from knowledge of the preferred antisense targets for the modulation of the synthesis of inflammatory cell adhesion molecules.

Several preferred embodiments of this invention are exemplified in accordance with the following nonlimiting examples. The target mRNA species for modulation relates to intercellular adhesion molecule-1, endothelial leukocyte adhesion molecule-1, and vascular cell adhesion molecule-1. Persons of ordinary skill in the art will appreciate that the present invention is not so limited, however, and that it is generally applicable. The inhibition or modulation of production of the ICAM-1 and/or ELAM-1 and/or VCAM-1 are expected to have significant therapeutic benefits in the treatment of disease. In order to assess the effectiveness of the compositions, an assay or series of assays is required.

- 17 -

EXAMPLES

Example 1

Expression of ICAM-1, VCAM-1 and ELAM-1 on the surface of cells can be quantitated using specific monoclonal antibodies in an ELISA. Cells are grown to confluence in 96 well microtiter plates. The cells are stimulated with either interleukin-1 or tumor necrosis factor for 4 to 8 hours to quantitate ELAM-1 and 8 to 24 hours to quantitate ICAM-1 and VCAM-1. Following the appropriate incubation time with the cytokine, the cells are gently washed three times with a buffered isotonic solution containing calcium and magnesium such as Dulbecco's phosphate buffered saline (D-PBS). The cells are then directly fixed on the microtiter plate with 1 to 2% paraformaldehyde diluted in D-PBS for 20 minutes at 25°C.

The cells are washed again with D-PBS three times. Nonspecific binding sites on the microtiter plate are blocked with 2% bovine serum albumin in D-PBS for 1 hour at 37°C. Cells are incubated with the appropriate monoclonal antibody diluted in blocking solution for 1 hour at 37°C. Unbound antibody is removed by washing the cells three times with D-PBS. Antibody bound to the cells is detected by incubation with a 1:1000 dilution of biotinylated goat anti-mouse IgG (Bethesda Research Laboratories, Gaithersburg, MD) in blocking solution for 1 hour at 37°C. Cells are washed three times with D-PBS and then incubated with a 1:1000 dilution of streptavidin conjugated to β -galactosidase (Bethesda Research Laboratories) for 1 hour at 37°C. The cells are washed three times with D-PBS for 5 minutes each. The amount of β -galactosidase bound to the specific monoclonal antibody is determined by developing the plate in a solution of 3.3 mM chlorophenolred- β -D-galactopyranoside, 50 mM sodium phosphate, 1.5 mM $MgCl_2$; pH=7.2 for 2 to 15 minutes at 37°C. The concentration of the product is determined by measuring the absorbance at 575 nm in an ELISA microtiter plate reader.

An example of the induction of ICAM-1 observed following stimulation with either interleukin-1 β or tumor necrosis factor α in several human cell lines is shown in

- 18 -

Figure 4. Cells were stimulated with increasing concentrations of interleukin-1 or tumor necrosis factor for 15 hours and processed as described above. ICAM-1 expression was determined by incubation with a 1:1000 dilution of the monoclonal antibody 84H10 (Amac Inc., Westbrook, ME). The cell lines used were passage 4 human umbilical vein endothelial cells (HUVEC), a human epidermal carcinoma cell line (A431), a human melanoma cell line (SK-MEL-2) and a human lung carcinoma cell line (A549). ICAM-1 was induced on all the cell lines, however, tumor necrosis factor was more effective than interleukin-1 in induction of ICAM-1 expression on the cell surface (Figure 4).

Screening antisense oligonucleotides for inhibition of ICAM-1, VCAM-1 or ELAM-1 expression is performed as described above with the exception of pretreatment of cells with the oligonucleotides prior to challenge with the cytokines. An example of antisense oligonucleotide inhibition of ICAM-1 expression is shown in Figure 5. Human umbilical vein endothelial cells (HUVEC) were treated with increasing concentration of oligonucleotide diluted in Opti MEM (GIBCO, Grand Island, NY) containing 8 μ M N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA) for 4 hours at 37°C to enhance uptake of the oligonucleotides. The medium was removed and replaced with endothelial growth medium (EGM-UV; Clonetics, San Diego, CA) containing the indicated concentration of oligonucleotide for an additional 4 hours. Interleukin-1 β was added to the cells at a concentration of 5 units/ml and incubated for 14 hours at 37°C. The cells were quantitated for ICAM-1 expression using a 1:1000 dilution of the monoclonal antibody 84H10 as described above. The oligonucleotides used were:

COMPOUND 1 - (ISIS 1558) a phosphodiester oligonucleotide designed to hybridize with position 64-80 of the mRNA covering the AUG initiation of translation codon having the sequence

5'-TGGGAGCCATAGCGAGGC-3' (SEQ ID NO: 1).

COMPOUND 2 - (ISIS 1570) a phosphorothioate containing oligonucleotide corresponding to the same sequence as COMPOUND

- 19 -

1.

COMPOUND 3 - a phosphorothioate oligonucleotide complementary to COMPOUND 1 and COMPOUND 2 exhibiting the sequence

5 5'-GCCTCGCTATGGCTCCCA-3' (SEQ ID NO: 81).

COMPOUND 4 - (ISIS 1572) a phosphorothioate containing oligonucleotide designed to hybridize to positions 2190-2210 of the mRNA in the 3' untranslated region containing the sequence

5'-GACACTCAATAAATAGCTGGT-3' (SEQ ID NO: 3).

10 COMPOUND 5 - (ISIS 1821) a phosphorothioate containing oligonucleotide designed to hybridize to human 5-lipoxygenase mRNA used as a control containing the sequence

5'-CATGGCGCGGGCCGCGGG-3' (SEQ ID NO: 82).

The phosphodiester oligonucleotide targeting the AUG
15 initiation of translation region of the human ICAM-1 mRNA (COMPOUND 1) did not inhibit expression of ICAM-1, however, the corresponding phosphorothioate containing oligonucleotide (COMPOUND 2) inhibited ICAM-1 expression by 70% at a concentration of 0.1 μ M and 90% at 1 μ M concentration (Figure
20 4). The increased potency of the phosphorothioate oligonucleotide over the phosphodiester is probably due to increased stability. The sense strand to COMPOUND 2, COMPOUND 3, modestly inhibited ICAM-1 expression at 10 μ M. If COMPOUND 2 was prehybridized to COMPOUND 3 prior to addition to the
25 cells, the effects of COMPOUND 2 on ICAM-1 expression were attenuated suggesting that the activity of COMPOUND 2 was due to antisense oligonucleotide effect, requiring hybridization to the mRNA. The antisense oligonucleotide directed against 3' untranslated sequences (COMPOUND 4) inhibited ICAM-1 expression
30 62% at a concentration of 1 μ M (Figure 5). The control oligonucleotide, targeting human 5-lipoxygenase (COMPOUND 5) reduced ICAM-1 expression by 20%. These data demonstrate that oligonucleotides are capable of inhibiting ICAM-1 expression on human umbilical vein endothelial cells and suggest that the
35 inhibition of ICAM-1 expression is due to an antisense activity.

The antisense oligonucleotide COMPOUND 2 at a

- 20 -

concentration of 1 μ M inhibits expression of ICAM-1 on human umbilical vein endothelial cells stimulated with increasing concentrations of tumor necrosis factor and interleukin-1 (Figure 6). These data demonstrate that the effects of COMPOUND 2 are not specific for interleukin-1 stimulation of cells.

Analogous assays can also be used to demonstrate inhibition of ELAM-1 and VCAM-1 expression by antisense oligonucleotides.

10 Example 2

A second cellular assay which can be used to demonstrate the effects of antisense oligonucleotides on ICAM-1, VCAM-1 or ELAM-1 expression is a cell adherence assay. Target cells are grown as a monolayer in a multiwell plate, treated with oligonucleotide followed by cytokine. The adhering cells are then added to the monolayer cells and incubated for 30 to 60 minutes at 37°C and washed to remove nonadhering cells. Cells adhering to the monolayer may be determined either by directly counting the adhering cells or prelabeled the cells with a radioisotope such as ^{51}Cr and quantitating the radioactivity associated with the monolayer as described. Dustin and Springer, *J. Cell Biol.* 1988, 107, 321-331. Antisense oligonucleotides may target either ICAM-1, VCAM-1 or ELAM-1 in the assay.

An example of the effects of antisense oligonucleotides targeting ICAM-1 mRNA on the adherence of DMSO differentiated HL-60 cells to tumor necrosis factor treated human umbilical vein endothelial cells is shown in Figure 7. Human umbilical vein endothelial cells were grown to 80% confluence in 12 well plates. The cells were treated with 2 μ M oligonucleotide diluted in Opti-MEM containing 8 μ M DOTMA for 4 hours at 37°C. The medium was removed and replaced with fresh endothelial cell growth medium (EGM-UV) containing 2 μ M of the indicated oligonucleotide and incubated 4 hours at 37°C. Tumor necrosis factor, 1 ng/ml, was added to cells as indicated and cells incubated for an additional 19 hours. The cells were

- 21 -

washed once with EGM-UV and 1.6×10^6 HL-60 cells differentiated for 4 days with 1.3% DMSO added. The cells were allowed to attach for 1 hour at 37°C and gently washed 4 times with Dulbecco's phosphate-buffered saline (D-PBS) warmed to 37°C. Adherent cells were detached from the monolayer by addition of 0.25 ml of cold (4°C) phosphate-buffered saline containing 5 mM EDTA and incubated on ice for 5 minutes. The number of cells removed by treatment with EDTA was determined by counting with a hemocytometer. Endothelial cells detached from the monolayer by EDTA treatment could easily be distinguished from HL-60 cells by morphological differences.

In the absence of tumor necrosis factor, 3% of the HL-60 cells bound to the endothelial cells. Treatment of the endothelial cell monolayer with 1 ng/ml tumor necrosis factor increased the number of adhering cells to 59% of total cells added (Figure 7). Treatment with the antisense oligonucleotide COMPOUND 2 or the control oligonucleotide COMPOUND 5 did not change the number of cells adhering to the monolayer in the absence of tumor necrosis factor treatment (Figure 7). The antisense oligonucleotide, COMPOUND 2 reduced the number of adhering cells from 59% of total cells added to 17% of the total cells added (Figure 7). In contrast, the control oligonucleotide COMPOUND 5 did not significantly reduce the number of cells adhering to the tumor necrosis factor treated endothelial monolayer, i.e., 53% of total cells added for COMPOUND 5 treated cells versus 59% for control cells.

These data indicate that antisense oligonucleotides are capable of inhibiting ICAM-1 expression on endothelial cells and that inhibition of ICAM-1 expression correlates with a decrease in the adherence of a neutrophil-like cell to the endothelial monolayer in a sequence specific fashion. Because other molecules also mediate adherence of white blood cells to endothelial cells, such as ELAM-1, and VCAM-1 it is not expected that adherence would be completely blocked.

Example 3

Synthesis and characterization of oligonucleotides

- 22 -

Unmodified DNA oligonucleotides were synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using standard phosphoramidite chemistry with oxidation by iodine. β -cyanoethyldiisopropyl-phosphoramidites were purchased from Applied Biosystems (Foster City, CA). For phosphorothioate oligonucleotides, the standard oxidation bottle was replaced by a 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation cycle wait step was increased to 68 seconds and was followed by the capping step.

2'-O-methyl phosphorothioate oligonucleotides were synthesized using 2'-O-methyl β -cyanoethyldiisopropyl-phosphoramidites (Chemgenes, Needham MA) and the standard cycle for unmodified oligonucleotides, except the wait step after pulse delivery of tetrazole and base was increased to 360 seconds. The 3'-base used to start the synthesis was a 2'-deoxyribonucleotide.

2'-fluoro phosphorothioate oligonucleotides were synthesized using 5'-dimethoxytrityl-3'-phosphoramidites and prepared as disclosed in U.S. patent application Serial No. 463,358, filed January 11, 1990, and 566,977, filed August 13, 1990, which are assigned to the same assignee as the instant application and which are incorporated by reference herein. The 2'-fluoro oligonucleotides were prepared using phosphoramidite chemistry and a slight modification of the standard DNA synthesis protocol: deprotection was effected using methanolic ammonia at room temperature.

After cleavage from the controlled pore glass column (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides were purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Analytical gel electrophoresis was accomplished in 20% acrylamide, 8 M urea, 45 mM Tris-borate buffer, pH 7.0. Oligodeoxynucleotides and phosphorothioate oligonucleotides were judged from electrophoresis to be greater than 80% full length material.

- 23 -

RNA oligonucleotide synthesis was performed on an ABI model 380B DNA synthesizer. The standard synthesis cycle was modified by increasing the wait step after the pulse delivery of tetrazole to 900 seconds. The bases were deprotected by incubation in methanolic ammonia overnight. Following base deprotections the oligonucleotides were dried in vacuo. The t-butyldimethylsilyl protecting the 2' hydroxyl was removed by incubating the oligonucleotide in 1 M tetrabutylammonium-fluoride in tetrahydrofuran overnight. The RNA oligonucleotides were further purified on C₁₈ Sep-Pak cartridges (Waters, Division of Millipore Corp., Milford MA) and ethanol precipitated.

The relative amounts of phosphorothioate and phosphodiester linkages obtained by this synthesis were periodically checked by ³¹P NMR spectroscopy. The spectra were obtained at ambient temperature using deuterium oxide or dimethyl sulfoxide-d₆ as solvent. Phosphorothioate samples typically contained less than one percent of phosphodiester linkages.

Secondary evaluation was performed with oligonucleotides purified by trityl-on HPLC on a PRP-1 column (Hamilton Co., Reno, Nevada) using a gradient of acetonitrile in 50 mM triethylammonium acetate, pH 7.0 (4% to 32% in 30 minutes, flow rate = 1.5 ml/min). Appropriate fractions were pooled, evaporated and treated with 5% acetic acid at ambient temperature for 15 minutes. The solution was extracted with an equal volume of ethyl acetate, neutralized with ammonium hydroxide, frozen and lyophilized. HPLC-purified oligonucleotides were not significantly different in potency from precipitated oligonucleotides, as judged by the ELISA assay for ICAM-1 expression.

- 24 -

Example 4**Cell culture and treatment with oligonucleotides**

The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (Bethesda MD). Cells were grown in Dulbecco's Modified Eagle's Medium (Irvine Scientific, Irvine CA) containing 1 gm glucose/liter and 10% fetal calf serum (Irvine Scientific). Human umbilical vein endothelial cells (HUVEC) (Clonetics, San Diego CA) were cultured in EGM-UV medium (Clonetics). HUVEC were used between the second and sixth passages. Human epidermal carcinoma A431 cells were obtained from the American Type Culture Collection and cultured in DMEM with 4.5 g/l glucose. Primary human keratinocytes were obtained from Clonetics and grown in KGM (Keratinocyte growth medium, Clonetics).

Cells grown in 96-well plates were washed three times with Opti-MEM (GIBCO, Grand Island, NY) prewarmed to 37°C. 100 µl of Opti-MEM containing either 10 µg/ml N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA, Bethesda Research Labs, Bethesda MD) in the case of HUVEC cells or 20 µg/ml DOTMA in the case of A549 cells was added to each well. Oligonucleotides were sterilized by centrifugation through 0.2 µm Centrex cellulose acetate filters (Schleicher and Schuell, Keene, NH). Oligonucleotides were added as 20x stock solution to the wells and incubated for 4 hours at 37°C. Medium was removed and replaced with 150 µl of the appropriate growth medium containing the indicated concentration of oligonucleotide. Cells were incubated for an additional 3 to 4 hours at 37°C then stimulated with the appropriate cytokine for 14 to 16 hours, as indicated. ICAM-1 expression was determined as described in Example 1. The presence of DOTMA during the first 4 hours incubation with oligonucleotide increased the potency of the oligonucleotides at least 100-fold. This increase in potency correlated with an increase in cell uptake of the oligonucleotide.

- 25 -

Example 5

**ELISA screening of additional antisense
oligonucleotides for activity against ICAM-1
gene expression in Interleukin-1 β -stimulated cells**

5 Antisense oligonucleotides were originally designed
that would hybridize to five target sites on the human ICAM-1
mRNA. Oligonucleotides were synthesized in both phosphodiester
(P=O; ISIS 1558, 1559, 1563, 1564 and 1565) and
phosphorothioate (P=S; ISIS 1570, 1571, 1572, 1573, and 1574)
10 forms. The oligonucleotides are shown in Table 1.

- 26 -

TABLE 1
 ANTISENSE OLIGONUCLEOTIDES WHICH TARGET HUMAN ICAM-1

| <u>ISIS NO.</u> | <u>SEQ ID NO.</u> | <u>TARGET REGION</u> | <u>MODIFICATION</u> |
|-----------------|-------------------|-------------------------------|---------------------|
| 1558 | 1 | AUG Codon (64-81) | P=O |
| 5 1559 | 2 | 5'-Untranslated (32-49) | P=O |
| 1563 | 3 | 3'-Untranslated (2190-3010) | P=O |
| 1564 | 4 | 3'-Untranslated (2849-2866) | P=O |
| 1565 | 5 | Coding Region (1378-1395) | P=O |
| 1570 | 1 | AUG Codon (64-81) | P=S |
| 10 1571 | 2 | 5'-Untranslated (32-49) | P=S |
| 1572 | 3 | 3'-Untranslated (2190-3010) | P=S |
| 1573 | 4 | 3'-Untranslated (2849-2866) | P=S |
| 1574 | 5 | Coding Region (1378-1395) | P=S |
| 1930 | 6 | 5'-Untranslated (1-20) | P=S |
| 15 1931 | 7 | AUG Codon (55-74) | P=S |
| 1932 | 8 | AUG Codon (72-91) | P=S |
| 1933 | 9 | Coding Region (111-130) | P=S |
| 1934 | 10 | Coding Region (351-370) | P=S |
| 1935 | 11 | Coding Region (889-908) | P=S |
| 20 1936 | 12 | Coding Region (1459-1468) | P=S |
| 1937 | 13 | Termination Codon (1651-1687) | P=S |
| 1938 | 14 | Termination Codon (1668-1687) | P=S |
| 1939 | 15 | 3'-Untranslated (1952-1971) | P=S |
| 1940 | 16 | 3'-Untranslated (2975-2994) | P=S |
| 25 2149 | 17 | AUG Codon (64-77) | P=S |
| 2163 | 18 | AUG Codon (64-75) | P=S |
| 2164 | 19 | AUG Codon (64-73) | P=S |
| 2165 | 20 | AUG Codon (66-80) | P=S |

- 27 -

| | | | | |
|----|------|----|-----------------------------|---|
| | 2173 | 21 | AUG Codon (64-79) | P=S |
| | 2302 | 22 | 3'-Untranslated (2114-2133) | P=S |
| | 2303 | 23 | 3'-Untranslated (2039-2058) | P=S |
| | 2304 | 24 | 3'-Untranslated (1895-1914) | P=S |
| 5 | 2305 | 25 | 3'-Untranslated (1935-1954) | P=S |
| | 2307 | 26 | 3'-Untranslated (1976-1995) | P=S |
| | 2634 | 1 | AUG-Codon (64-81) | 2'-fluoro A,C & U; P=S |
| 10 | 2637 | 15 | 3'-Untranslated (1952-1971) | 2'-fluoro A, C & U; |
| | 2691 | 1 | AUG Codon (64-81) | P=O, except last 3 bases, P=S |
| 15 | 2710 | 15 | 3'-Untranslated (1952-1971) | 2' - O - methyl; P=O |
| | 2711 | 1 | AUG Codon (64-81) | 2' - O - methyl; P=O |
| 20 | 2973 | 15 | 3'-Untranslated (1952-1971) | 2' - O - methyl; P=S |
| | 2974 | 1 | AUG Codon (64-81) | 2' - O - methyl; P=S |
| 25 | 3064 | 27 | 5'-CAP (32-51) | P=S; G & C added as spacer to 3' |
| | 3067 | 84 | 5'-CAP (32-51) | P=S |
| | 3222 | 84 | 5'-CAP (32-51) | 2' - O - methyl; P=O |
| 30 | 3224 | 84 | 5'-CAP (32-51) | 2' - O - methyl; P=S |
| | 3581 | 85 | 3'-Untranslated (1959-1978) | P=S |

Inhibition of ICAM-1 expression on the surface of interleukin-1 β -stimulated cells by the oligonucleotides was determined by
 35 ELISA assay as described in Example 1. The oligonucleotides
 were tested in two different cell lines. None of the

- 28 -

phosphodiester oligonucleotides inhibited ICAM-1 expression. This is probably due to the rapid degradation of phosphodiester oligonucleotides in cells. Of the five phosphorothioate oligonucleotides, the most active was ISIS 1570, which
5 hybridizes to the AUG translation initiation codon. A 2'-O-methyl phosphorothioate oligonucleotide, ISIS 2974, was approximately threefold less effective than ISIS 1570 in inhibiting ICAM-1 expression in HUVEC and A549 cells. A 2'-fluoro oligonucleotide, ISIS 2634, was also less effective.

10 Based on the initial data obtained with the five original targets, additional oligonucleotides were designed which would hybridize with the ICAM-1 mRNA. The antisense oligonucleotide (ISIS 3067) which hybridizes to the predicted transcription initiation site (5' cap site) was approximately as active in
15 IL-1 β -stimulated cells as the oligonucleotide that hybridizes to the AUG codon (ISIS 1570), as shown in Figure 8. ISIS 1931 and 1932 hybridize 5' and 3', respectively, to the AUG translation initiation codon. All three oligonucleotides that hybridize to the AUG region inhibit ICAM-1 expression, though
20 ISIS 1932 was slightly less active than ISIS 1570 and ISIS 1931. Oligonucleotides which hybridize to the coding region of ICAM-1 mRNA (ISIS 1933, 1934, 1935, 1574 and 1936) exhibited weak activity. Oligonucleotides that hybridize to the translation termination codon (ISIS 1937 and 1938) exhibited
25 moderate activity.

Surprisingly, the most active antisense oligonucleotide was ISIS 1939, a phosphorothioate oligonucleotide targeted to a sequence in the 3'- untranslated region of ICAM-1 mRNA (see Table 1). Other oligonucleotides having the same sequence were
30 tested, 2'-O-methyl (ISIS 2973) and 2'-fluoro (ISIS 2637); however, they did not exhibit this level of activity. Oligonucleotides targeted to other 3' untranslated sequences (ISIS 1572, 1573 and 1940) were also not as active as ISIS-1939. In fact, ISIS 1940, targeted to the polyadenylation
35 signal, did not inhibit ICAM-1 expression.

Because ISIS 1939 proved unexpectedly to exhibit the greatest antisense activity of the original 16 oligonucleotides

- 29 -

tested, other oligonucleotides were designed to hybridize to sequences in the 3'-untranslated region of ICAM-1 mRNA (ISIS 2302, 2303, 2304, 2305, and 2307, as shown in Table 1). ISIS 2307, which hybridizes to a site only five bases 3' to the ISIS 1939 target, was the least active of the series (Figure 8). ISIS 2302, which hybridizes to the ICAM-1 mRNA at a position 143 bases 3' to the ISIS 1939 target, was the most active of the series, with activity comparable to that of ISIS 1939. Examination of the predicted RNA secondary structure of the human ICAM-1 mRNA 3'-untranslated region (according to M. Zuker, *Science* 1989, 244, 48-52) revealed that both ISIS 1939 and ISIS 2302 hybridize to sequences predicted to be in a stable stem-loop structure. Current dogma suggests that regions of RNA secondary structure should be avoided when designing antisense oligonucleotides. Thus, ISIS 1939 and ISIS 2302 would not have been predicted to inhibit ICAM-1 expression.

The control oligonucleotide ISIS 1821 did inhibit ICAM-1 expression in HUVEC cells with activity comparable to that of ISIS 1934; however, in A549 cells ISIS 1821 was less effective than ISIS 1934. The negative control, ISIS 1821, was found to have a small amount of activity against ICAM expression, probably due in part to its ability to hybridize (12 of 13 base match) to the ICAM-1 mRNA at a position 15 bases 3' to the AUG translation initiation codon.

These studies indicate that the AUG translation initiation codon and specific 3'-untranslated sequences in the ICAM-1 mRNA were the most susceptible to antisense oligonucleotide inhibition of ICAM-1 expression.

In addition to inhibiting ICAM-1 expression in human umbilical vein cells and the human lung carcinoma cells (A549), ISIS 1570, ISIS 1939 and ISIS 2302 were shown to inhibit ICAM-1 expression in the human epidermal carcinoma A431 cells and in primary human keratinocytes (shown in Figure 9). These data demonstrate that antisense oligonucleotides are capable of inhibiting ICAM-1 expression in several human cell lines. Furthermore, the rank order potency of the oligonucleotides is

- 30 -

the same in the four cell lines examined. The fact that ICAM-1 expression could be inhibited in primary human keratinocytes is important because epidermal keratinocytes are an intended target of the antisense nucleotides.

5 Example 6

Antisense oligonucleotide inhibition of ICAM-1 expression in cells stimulated with other cytokines

Two oligonucleotides, ISIS 1570 and ISIS 1939, were tested for their ability to inhibit TNF- α and IFN- γ -induced ICAM-1 expression. Treatment of A549 cells with 1 μ M antisense oligonucleotide inhibited IL-1 β , TNF- α and IFN- γ -induced ICAM-1 expression in a sequence-specific manner. The antisense oligonucleotides inhibited IL-1 β and TNF- α -induced ICAM-1 expression to a similar extent, while IFN- γ -induced ICAM-1 expression was more sensitive to antisense inhibition. The control oligonucleotide, ISIS 1821, did not significantly inhibit IL-1 β - or TNF- α -induced ICAM-1 expression and inhibited IFN- γ -induced ICAM-1 expression slightly, as follows:

| | | Antisense Oligonucleotide (% Control Expression) | | |
|-----------------|---------------|---|------------------|------------------|
| <u>Cytokine</u> | | <u>ISIS 1570</u> | <u>ISIS 1939</u> | <u>ISIS 1821</u> |
| 3 U/ml | IL-1 β | 56.6 \pm 2.9 | 38.1 \pm 3.2 | 95 \pm 6.6 |
| 1 ng/ml | TNF- α | 58.1 \pm 0.9 | 37.6 \pm 4.1 | 103.5 \pm 8.2 |
| 100 U/ml | gamma-IFN | 38.9 \pm 3.0 | 18.3 \pm 7.0 | 83.0 \pm 3.5 |

Example 7

Antisense effects are abolished by sense strand controls

The antisense oligonucleotide inhibition of ICAM-1 expression by the oligonucleotides ISIS 1570 and ISIS 1939 could be reversed by hybridization of the oligonucleotides with their respective sense strands. The phosphorothioate sense strand for ISIS 1570 (ISIS 1575), when applied alone, slightly enhanced IL-1 β -induced ICAM-1 expression. Premixing ISIS 1570 with ISIS 1575 at equal molar concentrations, prior to addition

- 31 -

to the cells, blocked the effects of ISIS 1570. The complement to ISIS 1939 (ISIS 2115) enhanced ICAM-1 expression by 46% when added to the cells alone. Prehybridization of ISIS 2115 to ISIS 1939 completely blocked the inhibition of ICAM-1 expression by ISIS 1939.

Example 8

Measurement of oligonucleotide T_m (dissociation temperature of oligonucleotide from target)

To determine if the potency of the inhibition of ICAM-1 expression by antisense oligonucleotides was due to their affinity for their target sites, thermodynamic measurements were made for each of the oligonucleotides. The antisense oligonucleotides (synthesized as phosphorothioates) were hybridized to their complementary DNA sequences (synthesized as phosphodiesteres). Absorbance vs. temperature profiles were measured at 4 μ M each strand oligonucleotide in 100 mM Na⁺, 10 mM phosphate, 0.1 mM EDTA, pH 7.0. T_m 's and free energies of duplex formation were obtained from fits of data to a two-state model with linear sloping baselines (Petersheim, M. and D.H. Turner, *Biochemistry* 1983, 22, 256-263). Results are averages of at least three experiments.

When the antisense oligonucleotides were hybridized to their complementary DNA sequences (synthesized as phosphodiesteres), all of the antisense oligonucleotides with the exception of ISIS 1940 exhibited a T_m of at least 50°C. All the oligonucleotides should therefore be capable of hybridizing to the target ICAM-1 mRNA if the target sequences were exposed. Surprisingly, the potency of the antisense oligonucleotide did not correlate directly with either T_m or ΔG°_{37} . The oligonucleotide with the greatest biological activity, ISIS 1939, exhibited a T_m which was lower than that of the majority of the other oligonucleotides. Thus, hybridization affinity is not sufficient to ensure biological activity.

Example 9

Effect of oligonucleotide length on

- 32 -

antisense inhibition of ICAM-1 expression

The effect of oligonucleotide length on antisense activity was tested using truncated versions of ISIS 1570 (ISIS 2165, 2173, 2149, 2163 and 2164) and ISIS 1939 (ISIS 2540, 2544, 2545, 2546, 2547 and 2548). In general, antisense activity decreased as the length of the oligonucleotides decreased. Oligonucleotides 16 bases in length exhibited activity slightly less than 18 base oligonucleotides. Oligonucleotides 14 bases in length exhibited significantly less activity, and oligonucleotides 12 or 10 bases in length exhibited only weak activity. Examination of the relationship between oligonucleotide length and T_m and antisense activity reveals that a sharp transition occurs between 14 and 16 bases in length, while T_m increases linearly with length (Figure 10).

Example 10**Specificity of antisense inhibition of ICAM-1**

The specificity of the antisense oligonucleotides ISIS 1570 and ISIS 1939 for ICAM-1 was evaluated by immunoprecipitation of ^{35}S -labelled proteins. A549 cells were grown to confluence in 25 cm² tissue culture flasks and treated with antisense oligonucleotides as described in Example 4. The cells were stimulated with interleukin-1 β for 14 hours, washed with methionine-free DMEM plus 10% dialyzed fetal calf serum, and incubated for 1 hour in methionine-free medium containing 10% dialyzed fetal calf serum, 1 μM oligonucleotide and interleukin-1 β as indicated. ^{35}S -Methionine/cysteine mixture (Tran ^{35}S -label, purchased from ICN, Costa Mesa, CA) was added to the cells to an activity of 100 $\mu\text{Ci/ml}$ and the cells were incubated an additional 2 hours. Cellular proteins were extracted by incubation with 50 mM Tris-HCl pH 8.0, 150 mM NaCl, 1.0% NP-40, 0.5% deoxycholate and 2 mM EDTA (0.5 ml per well) at 4°C for 30 minutes. The extracts were clarified by centrifugation at 18,000 $\times g$ for 20 minutes. The supernatants were preadsorbed with 200 μl protein G-Sepharose beads (Bethesda Research Labs, Bethesda MD) for 2 hours at 4°C,

- 33 -

divided equally and incubated with either 5 µg ICAM-1 monoclonal antibody (purchased from AMAC Inc., Westbrook ME) or HLA-A,B antibody (W6/32, produced by murine hybridoma cells obtained from the American Type Culture Collection, Bethesda, MD) for 15 hours at 4°C. Immune complexes were trapped by incubation with 200 µl of a 50% suspension of protein G-Sepharose (v/v) for 2 hours at 4°C, washed 5 times with lysis buffer and resolved on an SDS-polyacrylamide gel. Proteins were detected by autoradiography.

10 Treatment of A549 cells with 5 units/ml of interleukin-1β was shown to result in the synthesis of a 95-100 kDa protein migrating as a doublet which was immunoprecipitated with the monoclonal antibody to ICAM-1. The appearance as a doublet is believed to be due to differently glycosylated forms of ICAM-1. Pretreatment of the cells with the antisense oligonucleotide ISIS 1570 at a concentration of 1 µM decreased the synthesis of ICAM-1 by approximately 50%, while 1 µM ISIS 1939 decreased ICAM-1 synthesis to near background. Antisense oligonucleotide ISIS 1940, inactive in the ICAM-1 ELISA assay

15 (Examples 1 and 5) did not significantly reduce ICAM-1 synthesis. None of the antisense oligonucleotides hybridizable with ICAM-1 targets had a demonstrable effect on HLA-A, B synthesis, demonstrating the specificity of the oligonucleotides for ICAM-1. Furthermore, the proteins which

20 nonspecifically precipitated with the ICAM-1 antibody and protein G-Sepharose were not significantly affected by treatment with the antisense oligonucleotides.

25

Example 11

30 Screening of additional antisense oligonucleotides for activity against ICAM-1 by cell adhesion assay

Human umbilical vein endothelial (HUVEC) cells were grown and treated with oligonucleotides as in Example 4. Cells were treated with either ISIS 1939, ISIS 1940, or the control oligonucleotide ISIS 1821 for 4 hours, then stimulated with

35 TNF-α for 20 hours. Basal HUVEC minimally bound HL-60 cells, while TNF-stimulated HUVEC bound 19% of the total cells added. Pretreatment of the HUVEC monolayer with 0.3 µM ISIS 1939

- 34 -

reduced the adherence of HL-60 cells to basal levels, as shown in Figure 11. The control oligonucleotide, ISIS 1821, and ISIS 1940 reduced the percentage of cells adhering from 19% to 9%. These data indicate that antisense oligonucleotides targeting
5 ICAM-1 may specifically decrease adherence of a leukocyte-like cell line (HL-60) to TNF- α -treated HUVEC.

Example 12

ELISA screening of antisense oligonucleotides for activity against ELAM-1 gene expression

10 Primary human umbilical vein endothelial (HUVEC) cells, passage 2 to 5, were plated in 96-well plates and allowed to reach confluence. Cells were washed three times with Opti-MEM (GIBCO, Grand Island NY). Cells were treated with increasing concentrations of oligonucleotide diluted in
15 Opti-MEM containing 10 μ g/ml DOTMA solution (Bethesda Research Labs, Bethesda MD) for 4 hours at 37°C. The medium was removed and replaced with EGM-UV (Clonetics, San Diego CA) plus oligonucleotide. Tumor necrosis factor α was added to the medium (2.5 ng/ml) and the cells were incubated an additional
20 4 hours at 37°C.

ELAM-1 expression was determined by ELISA. Cells were gently washed three times with Dulbecco's phosphate-buffered saline (D-PBS) prewarmed to 37°C. Cells were fixed with 95% ethanol at 4°C for 20 minutes, washed three times with D-PBS
25 and blocked with 2% BSA in D-PBS. Cells were incubated with ELAM-1 monoclonal antibody BBA-1 (R&D Systems, Minneapolis MN) diluted to 0.5 μ g/ml in D-PBS containing 2% BSA for 1 hour at 37°C. Cells were washed three times with D-PBS and the bound ELAM-1 antibody detected with biotinylated goat anti-mouse
30 secondary antibody followed by B-galactosidase-conjugated streptavidin as described in Example 1.

The activity of antisense phosphorothioate oligonucleotides which target 11 different regions on the ELAM-1 cDNA and two oligonucleotides which target ICAM-1 (as
35 controls) was determined using the ELAM-1 ELISA. The oligonucleotide and targets are shown in Table 2.

- 35 -

TABLE 2

ANTISENSE OLIGONUCLEOTIDES WHICH TARGET HUMAN ELAM-1

| | <u>ISIS NO.</u> | <u>SEQ ID NO.</u> | <u>TARGET REGION</u> | <u>MODIFICATION</u> |
|----|-----------------|-------------------|-----------------------------|---------------------|
| | 1926 | 28 | AUG Codon (143-164) | P=S |
| 5 | 2670 | 29 | 3'-Untranslated (3718-3737) | P=S |
| | 2673 | 30 | 3'-Untranslated (2657-2677) | P=S |
| | 2674 | 31 | 3'-Untranslated (2617-2637) | P=S |
| | 2678 | 32 | 3'-Untranslated (3558-3577) | P=S |
| | 2679 | 33 | 5'-Untranslated (41-60) | P=S |
| 10 | 2680 | 34 | 3'-Untranslated (3715-3729) | P=S |
| | 2683 | 35 | AUG Codon (143-163) | P=S |
| | 2686 | 36 | AUG Codon (149-169) | P=S |
| | 2687 | 37 | 5'-Untranslated (18-37) | P=S |
| | 2693 | 38 | 3'-Untranslated (2760-2788) | P=S |
| 15 | 2694 | 39 | 3'-Untranslated (2934-2954) | P=S |

In contrast to what was observed with antisense oligonucleotides targeted to ICAM-1 (Example 5), the most potent oligonucleotide modulator of ELAM-1 activity (ISIS 2679) was hybridizable with specific sequences in the 5'-untranslated region of ELAM-1. ISIS 2687, an oligonucleotide which hybridized to sequences ending three bases upstream of the ISIS 2679 target, did not show significant activity (Figure 12). Therefore, ISIS 2679 hybridizes to a unique site on the ELAM-1 mRNA, which is uniquely sensitive to inhibition with antisense oligonucleotides. The sensitivity of this site to inhibition with antisense oligonucleotides was not predictable based upon RNA secondary structure predictions or information in the literature.

Example 13

30 ELISA screening of additional antisense oligonucleotides for activity against ELAM-1 gene expression

- 36 -

Inhibition of ELAM-1 expression by eighteen antisense phosphorothioate oligonucleotides was determined using the ELISA assay as described in Example 12. The target sites of these oligonucleotides on the ELAM-1 mRNA are shown in Figure 5 13. The sequence and activity of each oligonucleotide against ELAM-1 are shown in Table 3. The oligonucleotides indicated by an asterisk (*) have IC₅₀'s of approximately 50 nM or below and are preferred. IC₅₀ indicates the dosage of oligonucleotide which results in 50% inhibition of ELAM-1 expression.

TABLE 3

Inhibition of human ELAM-1 expression by antisense oligonucleotides
ELAM-1 expression is given as % of control

| ISIS# | SEQ ID NO: | POSITION | SEQUENCE | VCAM-1 EXPRESSION | |
|-------|------------|--------------------|-----------------------|-------------------|-------------|
| | | | | 30 nM oligo | 50 nM oligo |
| 5 | *4764 | 5'-UTR 1-19 | GAAGTCAGCCCAAGAACAGCT | 70.2 | 50.2 |
| | 2687 | 5'-UTR 17-36 | TATAGGAGTTTGTGATGAA | 91.1 | 73.8 |
| | *2679 | 5'-UTR 40-59 | CTGCTGCCCTCTGTCTCAGGT | 6.4 | 6.0 |
| | *4759 | 5'-UTR 64-83 | ACAGGATCTCTCAGGTGGGT | 30.0 | 20.2 |
| | *2683 | AUG 143-163 | AATCATGACTTCAAGAGTTCT | 47.9 | 48.5 |
| 10 | *2686 | AUG 148-168 | TGAAGCAATCATGACTTCAAG | 51.1 | 46.9 |
| | *4756 | I/E 177-196 | CCAAAGTGAGAGCTGAGAGA | 53.9 | 35.7 |
| | 4732 | Coding 1936-1955 | CTGATTCAAGGCTTTGGCAG | 68.5 | 55.3 |
| | *4730 | I/E 3'UTR2006-2025 | TTCCCCAGATGCACCTGTTT | 14.1 | 2.3 |
| | *4729 | 3'-UTR 2063-2082 | GGGCCAGAGACCCGAGGAGA | 49.4 | 46.3 |
| 15 | *2674 | 3'-UTR 2617-2637 | CACAATCCTTAAGAACTCTTT | 33.5 | 28.1 |
| | 2673 | 3'-UTR 2656-2676 | GTATGGAAGATTATAATATAT | 58.9 | 53.8 |
| | 2694 | 3'-UTR 2933-2953 | GACAATATACAAACCTTCCAT | 72.0 | 64.6 |
| | *4719 | 3'-UTR 2993-3012 | ACGTTTGGCCTCATGGAAGT | 36.8 | 34.7 |
| | 4720 | 3'-UTR 3093-3112 | GGAATGCAAAAGCACATCCAT | 63.5 | 70.6 |
| 20 | *2678 | 3'-UTR 3557-3576 | ACCTCTGCTGTTCTGATCCT | 24.9 | 15.3 |
| | 2670 | 3'-UTR 3717-3736 | ACCACACTGGTATTTCACAC | 72.2 | 67.2 |

I/E indicates Intron/Exon junction

Oligonucleotides with IC50's of approximately 50 nM or below are indicated by an asterisk (*).

- 38 -

An additional oligonucleotide targeted to the 3'-untranslated region (ISIS 4728) did not inhibit ELAM expression.

Example 14

5 ELISA screening of antisense oligonucleotides for activity against VCAM-1 gene expression

Inhibition of VCAM-1 expression by fifteen antisense phosphorothioate oligonucleotides was determined using the ELISA assay approximately as described in Example 12, except that cells were stimulated with TNF- α for 16 hours and VCAM-1
10 expression was detected by a VCAM-1 specific monoclonal antibody (R & D Systems, Minneapolis, MN) used at 0.5 μ g/ml. The target sites of these oligonucleotides on the VCAM-1 mRNA are shown in Figure 14. The sequence and activity of each oligonucleotide against VCAM-1 are shown in Table 4. The
15 oligonucleotides indicated by an asterisk (*) have IC₅₀'s of approximately 50 nM or below and are preferred. IC₅₀ indicates the dosage of oligonucleotide which results in 50% inhibition of VCAM-1 expression.

TABLE 4

Inhibition of human VCAM-1 expression by antisense oligonucleotides
VCAM-1 expression is given as % of control

| 5 | ISIS# | SEQ ID NO: | POSITION | SEQUENCE | VCAM-1 EXPRESSION | |
|----|-------|---------------|----------|-----------------------|-------------------|-------------|
| | | | | | 30 nM oligo | 50 nM oligo |
| 5 | *5884 | 60 | 5'-UTR | CGATGCAGATACCGCGGAGT | 79.2 | 37.2 |
| | 3791 | 61 | 5'-UTR | GCCTGGGAGGGTATTCAGCT | 92.6 | 58.0 |
| | 5862 | 62 | 5'-UTR | CCTGTGTGTCCTGGGAGGG | 115.0 | 83.5 |
| | *3792 | 63 | AUG | GGCATTTTAAGTTGCTGTCG | 68.7 | 33.7 |
| | 5863 | 64 | CODING | CAGCCTGCCTTACTGTGGGC | 95.8 | 66.7 |
| 10 | *5874 | 65 | CODING | CTTGAACAATTAATTCACCT | 66.5 | 35.3 |
| | 5885 | 66 | E/I | TTACCATTGACATAAAGTGTT | 84.4 | 52.4 |
| | *5876 | 67 | CODING | CTGTGTCCTCTGTCTCCGCT | 43.5 | 26.6 |
| | *5875 | 68 | CODING | GTCTTTGTTGTTTCTCTTCC | 59.2 | 34.8 |
| | 3794 | 69 | TERMIN. | TGAACATATCAAGCATTAGC | 75.3 | 52.6 |
| 15 | *3800 | 70 | 3'-UTR | GCAATCTTGCTATGGCATAA | 64.4 | 47.7 |
| | *3805 | 71 | 3'-UTR | CCCGGCATCTTTACAAAACC | 67.7 | 44.9 |
| | *3801 | 50 | 3'-UTR | AACCCAGTGCTCCCTTTGCT | 36.5 | 21.3 |
| | *5847 | 72 | 3'-UTR | AACATCTCCGTACCATGCCA | 51.8 | 24.6 |
| | *3804 | 51 | 3'-UTR | GGCCACATTGGGAAAGTTGC | 55.1 | 29.3 |

E/I indicates exon/intron junction

Oligonucleotides with IC50's of approximately 50 nM or below are indicated by an asterisk (*).

- 40 -

Example 15

ICAM-1 expression in C8161 human melanoma cells: Human melanoma cell line C8161 (a gift of Dr. Dan Welch, Hershey Medical Center) was derived from an abdominal wall metastasis from a patient with recurrent malignant melanoma. These cells form multiple metastases in lung, subcutis, spleen, liver and regional lymph nodes after subcutaneous, intradermal and intravenous injection into athymic nude mice. Cells were grown in DMA-F12 medium containing 10% fetal calf serum and were passaged using 2 mM EDTA.

Exposure of C8161 cells to TNF- α resulted in a six-fold increase in cell surface expression of ICAM-1 and an increase in ICAM-1 mRNA levels in these cells. ICAM-1 expression on the cell surface was measured by ELISA. Cells were treated with increasing concentrations of antisense oligonucleotides in the presence of 15 μ g/ml Lipofectin for 4 hours at 37°C. ICAM-1 expression was induced by incubation with 5 ng/ml TNF- α for 16 hours. Cells were washed 3x in DPBS and fixed for 20 minutes in 2% formaldehyde. Cells were washed in DPBS, blocked with 2% BSA for 1 hour at 37°C and incubated with ICAM-1 monoclonal antibody 84H10 (AMAC, Inc., Westbrooke, ME). Detection of bound antibody was determined by incubation with a biotinylated goat anti-mouse IgG followed by incubation with β -galactosidase-conjugated streptavidin and developed with chlorophenol red- β -D-galactopyranoside and quantified by absorbance at 575 nm. ICAM-1 mRNA levels were measured by Northern blot analysis.

Example 16

Oligonucleotide inhibition of ICAM-1 expression in C8161 human melanoma cells: As shown in Figure 15, antisense oligonucleotides ICAM 1570 (SEQ ID NO: 1), ISIS 1939 (SEQ ID NO: 15) and ISIS 2302 (SEQ ID NO: 22) targeted to ICAM-1 decreased cell surface expression of ICAM-1 (detected by ELISA as in Example 16). ISIS 1822, a negative control oligonucleotide complementary to 5-lipoxygenase, did not affect ICAM-1 expression. The data were expressed as percentage of control activity, calculated as follows: (ICAM-1 expression for

- 41 -

oligonucleotide-treated, cytokine-induced cells)-(basal ICAM-1 expression)/(ICAM-1 cytokine-induced expression)-(basal ICAM-1 expression) x 100.

ISIS 1939 (SEQ ID NO: 15) and ISIS 2302 (SEQ ID NO: 22) markedly reduced ICAM-1 mRNA levels (detected by Northern blot analysis), but ISIS-1570 (SEQ ID NO: 1) decreased ICAM-1 mRNA levels only slightly.

Example 17

Experimental metastasis assay: To evaluate the role of ICAM-1 in metastasis, experimental metastasis assays were performed by injecting 1×10^5 C8161 cells into the lateral tail vein of athymic nude mice. Treatment of C8161 cells with the cytokine TNF- α and interferon γ has previously been shown to result in an increased number of lung metastases when cells were injected into nude mice [Miller, D.E. and Welch, D.R. (1990) Proc. Am. Assoc. Cancer Res. 13: 353].

After 4 weeks, mice were sacrificed, organs were fixed in Bouin's fixative and metastatic lesions on lungs were scored with the aid of a dissecting microscope. Four-week-old female athymic nude mice (Harlan Sprague Dawley) were used. Animals were maintained under the guidelines of the NIH. Groups of 4-8 mice each were tested in experimental metastasis assays.

Example 18

Antisense oligonucleotides ISIS 1570 and ISIS 2302 decrease metastatic potential of C8161 cells: Treatment of C8161 cells with antisense oligonucleotides ISIS 1570 and ISIS 2302, complementary to ICAM-1, was performed in the presence of the cationic lipid, Lipofectin (Gibco/BRL, Gaithersburg, MD). Antisense oligonucleotides were synthesized as described in Example 3. Cells were seeded in 60 mm tissue culture dishes at 10^6 cells/ml and incubated at 37°C for 3 days, washed with Opti-MEM (Gibco/BRL) 3 times and 100 μ l of Opti-MEM medium was added to each well. 0.5 μ M oligonucleotide and 15 μ g/ml lipofectin were mixed at room temperature for 15 minutes. 25 μ l

- 42 -

of the oligonucleotide-lipofectin mixture was added to the appropriate dishes and incubated at 37°C for 4 hr. The oligonucleotide-lipofectin mixture was removed and replaced with DME-F12 medium containing 10% fetal calf serum. After 4 hours, 500U/ml TNF- α was added to the appropriate wells and incubated for 18 hours at which time cells were removed from the plates, counted and injected into athymic nude mice.

Treatment of C8161 cells with ISIS 1570 (SEQ ID NO: 1) or ISIS 2302 (SEQ ID NO: 22) decreased the metastatic potential of these cells, and eliminated the enhanced metastatic ability of C8161 which resulted from TNF- α treatment. Data are shown in Table 5.

Table 5

Effect of antisense oligonucleotides to ICAM-1 on experimental metastasis of human melanoma cell line C8161

| Treatment | No. Lung Metastases per Mouse (Mean + S.E.M.) |
|--|--|
| Lipofectin only | 64 \pm 13 |
| Lipofectin + TNF- α | 81 \pm 8 |
| ISIS-1570 + Lipofectin | 38 \pm 15 |
| ISIS-2302 + Lipofectin | 23 \pm 6 |
| ISIS-1570 + Lipofectin + TNF- α | 49 \pm 6 |
| ISIS-2302 + Lipofectin + TNF- α | 31 \pm 8 |

| Treatment | No. Lung Metastases per Mouse (Mean + S.E.M.) |
|--|--|
| Lipofectin only | 64 \pm 13 |
| Lipofectin + TNF- α | 81 \pm 8 |
| ISIS-1570 + Lipofectin | 38 \pm 15 |
| ISIS-2302 + Lipofectin | 23 \pm 6 |
| ISIS-1570 + Lipofectin + TNF- α | 49 \pm 6 |
| ISIS-2302 + Lipofectin + TNF- α | 31 \pm 8 |

Example 19

Murine models for testing antisense oligonucleotides against ICAM-1: Many conditions which are believed to be mediated by intercellular adhesion molecules are not amenable to study in humans. For example, allograft rejection is a condition which is likely to be ameliorated by interference with ICAM-1 expression, but clearly this must be evaluated in animals rather than human transplant patients. Another such example is inflammatory bowel disease, and yet another is neutrophil migration (infiltration). These conditions can be tested in animal models, however, such as the mouse models used here.

Oligonucleotide sequences for inhibiting ICAM-1

- 43 -

expression in murine cells were identified. Murine ICAM-1 has approximately 50% homology with the human ICAM-1 sequence; a series of oligonucleotides which target the mouse ICAM-1 mRNA sequence were designed and synthesized, using information
5 gained from evaluation of oligonucleotides targeted to human ICAM-1. These oligonucleotides were screened for activity using an immunoprecipitation assay.

Murine DCEK-ICAM-1 cells (a gift from Dr. Adrienne Brian, University of California at San Diego) were treated with
10 1 μ M of oligonucleotide in the presence of 20 μ g/ml DOTMA/DOPE solution for 4 hours at 37°C. The medium was replaced with methionine-free medium plus 10% dialyzed fetal calf serum and 1 μ M antisense oligonucleotide. The cells were incubated for 1 hour in methionine-free medium, then 100 μ Ci/ml 35 S-labeled
15 methionine/cysteine mixture was added to the cells. Cells were incubated an additional 2 hours, washed 4 times with PBS, and extracted with buffer containing 20 mM Tris, pH 7.2, 20 mM KCl, 5 mM EDTA, 1% Triton X-100, 0.1 mM leupeptin, 10 μ g/ml aprotinin, and 1 mM PMSF. ICAM-1 was immunoprecipitated from
20 the extracts by incubating with a murine-specific ICAM-1 antibody (YN1/1.7.4) followed by protein G-sepharose. The immunoprecipitates were analyzed by SDS-PAGE and autoradiographed. Phosphorothioate oligonucleotides ISIS 3066 and 3069, which target the AUG codon of mouse ICAM-1, inhibited
25 ICAM-1 synthesis by 48% and 63%, respectively, while oligonucleotides ISIS 3065 and ISIS 3082, which target sequences in the 3'-untranslated region of murine ICAM-1 mRNA inhibited ICAM-1 synthesis by 47% and 97%, respectively. The most active antisense oligonucleotide against mouse ICAM-1 was
30 targeted to the 3'-untranslated region. ISIS 3082 was evaluated further based on these results; this 20-mer phosphorothioate oligonucleotide comprises the sequence (5' to 3') TGC ATC CCC CAG GCC ACC AT (SEQ ID NO: 83).

- 44 -

Example 20

Antisense oligonucleotides to ICAM-1 reduce inflammatory bowel disease in murine model system:

A mouse model for inflammatory bowel disease (IBD) has recently been developed. Okayasu et al., (1990) Gastroenterology 98:694-702. Administration of dextran sulfate to mice induces colitis that mimics human IBD in almost every detail. Dextran sulfate-induced IBD and human IBD have subsequently been closely compared at the histological level and the mouse model has been found to be an extremely reproducible and reliable model. It is used here to test the effect of ISIS 3082, a 20-base phosphorothioate antisense oligonucleotide which is complementary to the 3' untranslated region of the murine ICAM-1.

Female Swiss Webster mice (8 weeks of age) weighing approximately 25 to 30 grams are kept under standard conditions. Mice are allowed to acclimate for at least 5 days before initiation of experimental procedures. Mice are given 5% dextran sulfate sodium in their drinking water (available ad libitum) for 5 days. Concomitantly, ISIS 3082 oligonucleotide in pharmaceutical carrier, carrier alone (negative control) or TGF- β (known to protect against dextran sulfate-mediated colitis in mice) is administered. ISIS 3082 was given as daily subcutaneous injection of 1 mg/kg or 10 mg/kg for 5 days. TGF- β was given as 1 μ g/mouse intracolonicly. At 1 mg/kg, the oligonucleotide was as effective as TGF- β in protecting against dextran-sulfate-induced colitis.

Mice were sacrificed on day 6 and colons were subjected to histopathologic evaluation. Until sacrifice, disease activity was monitored by observing mice for weight changes and by observing stools for evidence of colitis. Mice were weighed daily. Stools were observed daily for changes in consistency and for presence of occult or gross bleeding. A scoring system was used to develop a disease activity index by which weight loss, stool consistency and presence of bleeding were graded on a scale of 0 to 3 (0 being normal and 3 being most severely affected) and an index was calculated. Drug-

- 45 -

induced changes in the disease activity index were analyzed statistically. The disease activity index has been shown to correlate extremely well with IBD in general. Results are shown in Figure 16. At 1 mg/kg, the oligonucleotide reduced the disease index by 40%.

Example 21

Antisense oligonucleotide to ICAM-1 increases survival in murine heterotopic heart transplant model: To determine the therapeutic effects of ICAM-1 antisense oligonucleotide in preventing allograft rejection the murine ICAM-1 specific oligonucleotide ISIS 3082 was tested for activity in a murine vascularized heterotopic heart transplant model. Hearts from Balb/c mice were transplanted into the abdominal cavity of C3H mice as primary vascularized grafts essentially as described by Isobe et al. [(1991) Circulation 84:1246-1255]. Oligonucleotides were administered by continuous intravenous administration via a 7-day Alzet pump. The mean survival time for untreated mice was 9.2 ± 0.8 days (8, 9, 9, 9, 10, 10 days). Treatment of the mice for 7 days with 5 mg/kg ISIS 3082 increased the mean survival time to 14.3 ± 4.6 days (11, 12, 13, 21 days).

Example 22

Antisense oligonucleotide to ICAM-1 decreases leukocyte migration: Leukocyte infiltration of tissues and organs is a major aspect of the inflammatory process and contributes to tissue damage resulting from inflammation. The effect of ISIS 3082 on leukocyte migration was examined using a mouse model in which carrageenan-soaked sponges were implanted subcutaneously. Carrageenan stimulates leukocyte migration and edema. Effect of oligonucleotide on leukocyte migration in inflammatory exudates is evaluated by quantitation of leukocytes infiltrating the implanted sponges. Following a four hour fast, 40 mice were assigned randomly to eight groups each containing five mice. Each mouse was anesthetized with Metofane® and a polyester sponge impregnated with 1 ml of a 20 mg/ml solution of carrageenan was implanted subcutaneously. Saline was administered intravenously

- 46 -

to Group 1 at 10 ml/kg four hours prior to sponge implantation and this served as the vehicle control. Indomethacin (positive control) was administered orally at 3 mg/kg at a volume of 20 ml/kg to Group 2 immediately following surgery, again 6-8 hours later and again at 21 hours post-implantation. ISIS 3082 was administered intravenously at 5 mg/kg to Group 3 four hours prior to sponge implantation. ISIS 3082 was administered intravenously at 5 mg/kg to Group 4 immediately following sponge implantation. ISIS 3082 was administered intravenously at 5 mg/kg to Groups 5, 6, 7 and 8 at 2, 4, 8 and 18 hours following sponge implantation, respectively. Twenty-four hours after implantation, sponges were removed, immersed in EDTA and saline (5 ml) and squeezed dry. Total numbers of leukocytes in sponge exudate mixtures were determined.

15 The oral administration of indomethacin at 3 mg/kg produced a 79% reduction in mean leukocyte count when compared to the vehicle control group.

 A 42% reduction in mean leukocyte count was observed following the administration of ISIS 3082 at 5 mg/kg four hours prior to sponge implantation (Group 3). A 47% reduction in mean leukocyte count was observed following the administration of ISIS 3082 at 5 mg/kg immediately following sponge implantation (Group 4). All animals appeared normal throughout the course of study.

- 47 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Bennett and Mirabelli

(ii) TITLE OF INVENTION: Oligonucleotide Modulation
of Cell Adhesion

(iii) NUMBER OF SEQUENCES: 85

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Woodland Falls Corporate Park

(B) STREET: 210 Lake Drive East, Suite 201

(C) CITY: Cherry Hill

(D) STATE: NJ

(E) COUNTRY: USA

(F) ZIP: 08002

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: DISKETTE, 3.5 INCH, 1.44 Mb STORAGE

(B) COMPUTER: IBM PS/2

(C) OPERATING SYSTEM: PC-DOS

(D) SOFTWARE: WORDPERFECT 5.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: n/a

(B) FILING DATE: Herewith

(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 939,855

(B) FILING DATE: September 2, 1992

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: PCT/US91/05209

(B) FILING DATE: July 23, 1991

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 567,286

(B) FILING DATE: August 14, 1990

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 007,997

(B) FILING DATE: January 20, 1993

- 48 -

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Jane Massey Licata
- (B) REGISTRATION NUMBER: 32,257
- (C) REFERENCE/DOCKET NUMBER: ISPH-0002

(ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: (609) 779-2400
- (B) TELEFAX: (609) 779-8488

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TGGGAGCCAT AGCGAGGC

18

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

GAGGAGCTCA GCGTCGACTG

20

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GACACTCAAT AAATAGCTGG T

21

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18

- 49 -

- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:
GAGGCTGAGG TGGGAGGA 18
- (2) INFORMATION FOR SEQ ID NO: 5:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:
CGATGGGCAG TGGGAAAG 18
- (2) INFORMATION FOR SEQ ID NO: 6:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:
GGGCGCGTGA TCCTTATAGC 20
- (2) INFORMATION FOR SEQ ID NO: 7:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:
CATAGCGAGG CTGAGGTTGC 20
- (2) INFORMATION FOR SEQ ID NO: 8:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid

- 50 -

- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:
CGGGGGCTGC TGGGAGCCAT 20
- (2) INFORMATION FOR SEQ ID NO: 9:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:
AGAGCCCCGA GCAGGACCAG 20
- (2) INFORMATION FOR SEQ ID NO: 10:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:
TGCCCATCAG GGCAGTTTGA 20
- (2) INFORMATION FOR SEQ ID NO: 11:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:
GGTCACACTG ACTGAGGCCT 20
- (2) INFORMATION FOR SEQ ID NO: 12:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single

- 51 -

- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:
CTCGCGGGTG ACCTCCCCTT 20
- (2) INFORMATION FOR SEQ ID NO: 13:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:
TCAGGGAGGC GTGGCTTGTG 20
- (2) INFORMATION FOR SEQ ID NO: 14:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:
CCTGTCCCGG GATAGGTTCA 20
- (2) INFORMATION FOR SEQ ID NO: 15:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:
CCCCCACCAC TTCCCCTCTC 20
- (2) INFORMATION FOR SEQ ID NO: 16:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- 52 -

- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:
TTGAGAAAGC TTTATTA ACT 20
- (2) INFORMATION FOR SEQ ID NO: 17:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 14
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:
AGCCATAGCG AGGC 14
- (2) INFORMATION FOR SEQ ID NO: 18:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 12
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:
CCATAGCGAG GC 12
- (2) INFORMATION FOR SEQ ID NO: 19:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 10
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:
ATAGCGAGGC 10
- (2) INFORMATION FOR SEQ ID NO: 20:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes

- 53 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:
TGGGAGCCAT AGCGAG 16
- (2) INFORMATION FOR SEQ ID NO: 21:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 16
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:
GGAGCCATAG CGAGGC 16
- (2) INFORMATION FOR SEQ ID NO: 22:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:
GCCCAAGCTG GCATCCGTCA 20
- (2) INFORMATION FOR SEQ ID NO: 23:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:
TCTGTAAGTC TGTGGGCCTC 20
- (2) INFORMATION FOR SEQ ID NO: 24:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

- 54 -

- AGTCTTGCTC CTCCTCTTG 20
- (2) INFORMATION FOR SEQ ID NO: 25:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:
- CTCATCAGGC TAGACTTTAA 20
- (2) INFORMATION FOR SEQ ID NO: 26:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:
- TGTCCTCATG GTGGGGCTAT 20
- (2) INFORMATION FOR SEQ ID NO: 27:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:
- TCTGAGTAGC AGAGGAGCTC GA 22
- (2) INFORMATION FOR SEQ ID NO: 28:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 22
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:
- CAATCATGAC TTCAAGAGTT CT 22

- 55 -

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

ACCACACTGG TATTTACAC

20

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

GTATGGAAGA TTATAATATA T

21

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

CACAATCCTT AAGAACTCTT T

21

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

ACCTCTGCTG TTCTGATCCT

20

(2) INFORMATION FOR SEQ ID NO: 33:

- 56 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:
 CTGCTGCCTC TGTCTCAGGT 20
- (2) INFORMATION FOR SEQ ID NO: 34:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:
 GGTATTGAC ACAGC 15
- (2) INFORMATION FOR SEQ ID NO: 35:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:
 AATCATGACT TCAAGAGTTC T 21
- (2) INFORMATION FOR SEQ ID NO: 36:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:
 TGAAGCAATC ATGACTTCAA G 21
- (2) INFORMATION FOR SEQ ID NO: 37:
(i) SEQUENCE CHARACTERISTICS:

- 57 -

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:
TATAGGAGTT TTGATGTGAA 20
- (2) INFORMATION FOR SEQ ID NO: 38:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:
ACAATGAGGG GGTAATCTAC A 21
- (2) INFORMATION FOR SEQ ID NO: 39:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:
GACAATATAC AAACCTTCCA T 21
- (2) INFORMATION FOR SEQ ID NO: 40:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:
CCAGGCATTT TAAGTTGCTG T 21
- (2) INFORMATION FOR SEQ ID NO: 41:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20

- 58 -

- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:
CCTGAAGCCA GTGAGGCCCG 20
- (2) INFORMATION FOR SEQ ID NO: 42:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:
GATGAGAAAA TAGTGAACC A 21
- (2) INFORMATION FOR SEQ ID NO: 43:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:
CTGAGCAAGA TATCTAGAT 19
- (2) INFORMATION FOR SEQ ID NO: 44:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:
CTACACTTTT GATTCTGT 19
- (2) INFORMATION FOR SEQ ID NO: 45:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22
 - (B) TYPE: Nucleic Acid

- 59 -

- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:
TTGAACATAT CAAGCATTAG CT 22
- (2) INFORMATION FOR SEQ ID NO: 46:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:
TTTACATATG TACAAATTAT GT 22
- (2) INFORMATION FOR SEQ ID NO: 47:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:
AATTATCACT TTACTATACA AA 22
- (2) INFORMATION FOR SEQ ID NO: 48:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:
AGGGCTGACC AAGACGGTTG T 21
- (2) INFORMATION FOR SEQ ID NO: 49:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single

- 60 -

- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:
CCATCTTCCC AGGCATTTTA 20
- (2) INFORMATION FOR SEQ ID NO: 50:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:
AACCCAGTGC TCCCTTTGCT 20
- (2) INFORMATION FOR SEQ ID NO: 51:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:
GGCCACATTG GGAAAGTTGC 20
- (2) INFORMATION FOR SEQ ID NO: 52:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:
GAAGTCAGCC AAGAACAGCT 20
- (2) INFORMATION FOR SEQ ID NO: 53:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

- 61 -

- (iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:
ACAGGATCTC TCAGGTGGGT 20
- (2) INFORMATION FOR SEQ ID NO: 54:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:
CCAAAGTGAG AGCTGAGAGA 20
- (2) INFORMATION FOR SEQ ID NO: 55:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:
CTGATTCAAG GCTTTGGCAG 20
- (2) INFORMATION FOR SEQ ID NO: 56:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:
TTCCCCAGAT GCACCTGTTT 20
- (2) INFORMATION FOR SEQ ID NO: 57:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes

- 62 -

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:
GGGCCAGAGA CCCGAGGAGA 20
- (2) INFORMATION FOR SEQ ID NO: 58:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:
ACGTTTGCC TCATGGAAGT 20
- (2) INFORMATION FOR SEQ ID NO: 59:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:
GGAATGCAAA GCACATCCAT 20
- (2) INFORMATION FOR SEQ ID NO: 60:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:
CGATGCAGAT ACCGCGGAGT 20
- (2) INFORMATION FOR SEQ ID NO: 61:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

- 63 -

- GCCTGGGAGG GTATTCAGCT 20
- (2) INFORMATION FOR SEQ ID NO: 62:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:
CCTGTGTGTG CCTGGGAGGG 20
- (2) INFORMATION FOR SEQ ID NO: 63:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:
GGCATTTTAA GTTGCTGTCG 20
- (2) INFORMATION FOR SEQ ID NO: 64:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:
CAGCCTGCCT TACTGTGGGC 20
- (2) INFORMATION FOR SEQ ID NO: 65:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:
CTTGAACAAT TAATTCCACC T 21

- 64 -

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

TTACCATTGA CATAAAGTGT T

21

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

CTGTGTCTCC TGTCTCCGCT

20

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

GTCTTTGTTG TTTTCTCTTCC

21

(2) INFORMATION FOR SEQ ID NO: 69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(iv) ANTI-SENSE: Yes

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

TGAACATATC AAGCATTAGC

20

(2) INFORMATION FOR SEQ ID NO: 70:

- 65 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:
GCAATCTTGC TATGGCATAA 20
- (2) INFORMATION FOR SEQ ID NO: 71:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
 (iv) ANTI-SENSE: Yes
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:
CCCGGCATCT TTACAAAACC 20
- (2) INFORMATION FOR SEQ ID NO: 72:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
 (iv) ANTI-SENSE: Yes
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:
AACATCTCCG TACCATGCCA 20
- (2) INFORMATION FOR SEQ ID NO: 73:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear
 (iv) ANTI-SENSE: Yes
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:
TCACTGCTGC CTCTGTCTCA GG 22
- (2) INFORMATION FOR SEQ ID NO: 74:
 (i) SEQUENCE CHARACTERISTICS:

- 66 -

- (A) LENGTH: 23
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:
TGATTCTTTT GAACTTAAAA GGA 23
- (2) INFORMATION FOR SEQ ID NO: 75:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:
TTAAAGGATG TAAGAAGGCT 20
- (2) INFORMATION FOR SEQ ID NO: 76:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:
CATAAGCACA TTTATTGTC 19
- (2) INFORMATION FOR SEQ ID NO: 77:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:
TTTTGGGAAG CAGTTGTTCA 20
- (2) INFORMATION FOR SEQ ID NO: 78:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21

- 67 -

- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear
- (iv) ANTI-SENSE: Yes
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:
AACTGTGAAG CAATCATGAC T 21
- (2) INFORMATION FOR SEQ ID NO: 79:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:
CCTTGAGTGG TGCATTCAAC CT 22
- (2) INFORMATION FOR SEQ ID NO: 80:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:
AATGCTTGCT CACACAGGCA TT 22
- (2) INFORMATION FOR SEQ ID NO: 81:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18
 - (B) TYPE: Nucleic Acid
 - (C) STRANDEDNESS: Single
 - (D) TOPOLOGY: Linear
 - (iv) ANTI-SENSE: Yes
 - (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:
GCCTCGCTAT GGCTCCCA 18
- (2) INFORMATION FOR SEQ ID NO: 82:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18
 - (B) TYPE: Nucleic Acid

- 68 -

- (C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:
CATGGCGCGG GCCGCGGG 18
- (2) INFORMATION FOR SEQ ID NO: 83:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:
TGCATCCCCC AGGCCACCAT 20
- (2) INFORMATION FOR SEQ ID NO: 84:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:
TCTGAGTAGC AGAGGAGCTC 20
- (2) INFORMATION FOR SEQ ID NO: 85:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: Nucleic Acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(iv) ANTI-SENSE: Yes
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:
TATGTCTCCC CCACCACTTC 20

- 69 -

What is claimed is:

1. An oligonucleotide specifically hybridizable with at least a portion of a nucleic acid encoding a protein capable of modulating cell adhesion.
- 5 2. The oligonucleotide of claim 1 which is specifically hybridizable with mRNA.
3. The oligonucleotide of claim 1 which is specifically hybridizable with a gene, forming a triple stranded structure for modulating the amount of mRNA made from said gene.
- 10 4. The oligonucleotide of claim 2 specifically hybridizable with a transcription initiation site, a translation initiation site, 5'-untranslated sequences, 3' untranslated sequences or an intron/exon junction.
5. The oligonucleotide of claim 4 specifically
15 hybridizable with 5' cap site or an intervening sequence.
6. The oligonucleotide of claim 1 wherein said protein is intercellular adhesion molecule-1.
7. The oligonucleotide of claim 1 wherein said protein is endothelial leukocyte adhesion molecule-1.
- 20 8. The oligonucleotide of claim 1 wherein said protein is vascular cell adhesion molecule-1.
9. The oligonucleotide of claim 1 comprising from about 3 to about 50 nucleotides.
10. The oligonucleotide of claim 1 comprising from
25 about 8 to about 25 nucleotides.
11. The oligonucleotide of claim 1 comprising from about 10 to about 20 nucleotides.

- 70 -

12. A pharmaceutical composition comprising an oligonucleotide of claim 1 and a pharmaceutically acceptable carrier.

13. The oligonucleotide of claim 1 wherein at least one of the linking groups between nucleotide units comprises a sulfur-containing species.

14. The oligonucleotide of claim 1 wherein at least one of the linking groups between nucleotide units comprises a phosphorothioate moiety.

10 15. An oligonucleotide comprising SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 13, 14, 15, 16, 22, 28, 40, 41, 42, 43, 44, 45, 46, 47, 48, 73, 74, 75, 76, 77, 78, 79, 80 or 85.

16. A pharmaceutical composition comprising an oligonucleotide of claim 15 and a pharmaceutically acceptable
15 carrier.

17. The oligonucleotide of claim 15 wherein at least one of the linking groups between nucleotide units comprises a sulfur containing species.

18. The oligonucleotide of claim 15 wherein at least
20 one of the linking groups between nucleotide units comprises a phosphorothioate moiety.

19. A method of modulating the synthesis of intercellular adhesion molecules in a cell or tissue comprising contacting the cell or tissue with an oligonucleotide
25 specifically hybridizable with nucleic acids encoding at least a portion of a protein capable of modulating cell adhesion and modulating the synthesis of intracellular adhesion molecules in the cell or tissue.

- 71 -

20. The method of claim 19 wherein the oligonucleotide is specifically hybridizable with a transcription initiation site, a translation initiation site, 5'-untranslated sequence, 3'-untranslated sequences, or an intervening sequence.
- 5 21. The method of claim 19 wherein said oligonucleotide is specifically hybridizable with a 5' cap site of mRNA or an adjacent sequence.
22. The method of claim 19 wherein said protein is intercellular adhesion molecule-1.
- 10 23. The method of claim 19 wherein said protein is endothelial leukocyte adhesion molecule-1.
24. The method of claim 19 wherein said protein is vascular cell adhesion molecule-1.
- 15 25. The method of claim 19 wherein the oligonucleotide comprises SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 13, 14, 15, 16, 22, 28, 40, 41, 42, 43, 44, 45, 46, 47, 48, 73, 74, 75, 76, 77, 78, 79, 80 or 85.
- 20 26. The method of claim 19 wherein at least one of the linking groups between nucleotide units comprises a sulfur-containing species.
27. The method of claim 19 wherein at least one of the linking groups between nucleotide units comprises a phosphorothioate moiety.

- 72 -

28. A method of treating an animal suspected of having a disease which is modulated by changes in intercellular adhesion molecules comprising contacting an animal with a therapeutically effective amount of an oligonucleotide specifically hybridizable
5 with nucleic acids encoding at least a portion of a protein which modulates the synthesis or metabolism of intercellular adhesion molecules.

29. The method of claim 28 wherein said oligonucleotide is specifically hybridizable to a transcription initiation site,
10 translation initiation site, 5'-untranslated sequences, 3' untranslated sequences or an intervening sequence.

30. The method of claim 28 wherein said oligonucleotide is specifically hybridizable with a 5' cap site of mRNA and adjacent sequences.

15 31. The method of claim 28 wherein said protein is intercellular adhesion molecule-1.

32. The method of claim 28 wherein said protein is endothelial leukocyte adhesion molecule-1.

33. The method of claim 28 wherein said protein is
20 vascular cell adhesion molecule-1.

34. The method of claim 28 wherein said oligonucleotide is administered in a pharmaceutical composition comprising the oligonucleotide and a pharmaceutically acceptable carrier.

35. The method of claim 28 wherein the oligonucleotide
25 is administered topically.

36. The method of claim 28 wherein the oligonucleotide is administered systemically.

- 73 -

37. The method of claim 28 wherein the oligonucleotide comprises SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8, 13, 14, 15, 16, 22, 28, 40, 41, 42, 43, 44, 45, 46, 47, 48, 73, 74, 75, 76, 77, 78, 79, 80 or 85.
- 5 38. The method of claim 28 wherein at least one of the linking groups between nucleotide units comprises a sulfur-containing species.
39. The method of claim 28 wherein at least one of the linking groups between nucleotide units comprises a
10 phosphorothioate moiety.
40. An oligonucleotide of claim 7 which is selected from the sequences in Table 3.
41. An oligonucleotide of claim 7 comprising: SEQ ID NO: 31, 32, 33, 35, 36, 52, 53, 54, 56, 57, or 58.
- 15 42. An oligonucleotide of claim 8 comprising a sequence identified in Table 4.
43. An oligonucleotide of claim 8 comprising SEQ ID NO: 50, 51, 60, 63, 65, 67, 68, 70, 71, or 72.
44. The method of claim 28 wherein said disease is
20 allograft rejection.
45. An oligonucleotide comprising SEQ ID NO: 1.
46. An oligonucleotide comprising SEQ ID NO: 15.
47. An oligonucleotide comprising SEQ ID NO: 22.
48. An oligonucleotide comprising SEQ ID NO: 33 or
25 56.

- 74 -

49. An oligonucleotide comprising SEQ ID NO: 50 or 67.
50. The method of claim 19 wherein the oligonucleotide comprises SEQ ID NO: 1, SEQ NO: 15, SEQ NO: 22 or SEQ ID NO: 84.
51. The method of claim 19 wherein the oligonucleotide comprises SEQ ID NO: 33 or 56.
52. The method of claim 19 wherein the oligonucleotide comprises SEQ ID NO: 50 or 67.
- 10 53. The method of claim 28 wherein the oligonucleotide comprises SEQ ID NO: 1, SEQ ID NO: 15, SEQ ID NO: 22 or SEQ ID NO: 84.
54. The method of claim 28 wherein the oligonucleotide comprises SEQ ID NO: 33 or 56.
- 15 55. The method of claim 28 wherein the oligonucleotide comprises SEQ ID NO: 50 or 67.
56. The method of claim 28 wherein said disease is psoriasis.
57. The method of claim 28 wherein said disease is 20 an inflammatory disease.
58. The method of claim 57 wherein said inflammatory disease is inflammatory bowel disease.
59. The method of claim 57 wherein said inflammatory disease is characterized by leukocyte infiltration.
- 25 60. The method of claim 28 wherein said disease is characterized by metastases.

- 75 -

61. The method of claim 60 wherein said metastases affect the lung.

62. The method of claim 28 wherein said disease is malignant melanoma.

5 63. A method of decreasing metastasis in an animal suspected of having a metastatic condition comprising contacting the animal with a therapeutically effective amount of an oligonucleotide specifically hybridizable with nucleic acids encoding at least a portion of a protein which modulates the
10 synthesis or metabolism of an intercellular adhesion molecule.

64. The method of claim 63 wherein said oligonucleotide is specifically hybridizable to a transcription initiation site, translation initiation site, 5'-untranslated sequences, 3' untranslated sequences or an intervening sequence.

15 65. The method of claim 63 wherein said oligonucleotide is specifically hybridizable with a 5' cap site of mRNA and adjacent sequences.

66. The method of claim 63 wherein said protein is intercellular adhesion molecule-1.

20 67. The method of claim 63 wherein said protein is endothelial leukocyte adhesion molecule-1.

68. The method of claim 63 wherein said protein is vascular cell adhesion molecule-1.

25 69. The method of claim 63 wherein said oligonucleotide is administered in a pharmaceutical composition comprising the oligonucleotide and a pharmaceutically acceptable carrier.

1/26

GCTATAAGGA TCACGGGCCC CAGTCGACGC TGAGCTCCTC TGCTACTCAG AGTTGCAACC TCAGCCTCGCT
 ATG GCT CCC AGC AGC CCC CGG CTG CCC GCA CTC CTG GTC CTG CTC GGG GCT CTG TTC CCA
 MET ALA PRO SER SER PRO ARG PRO ALA LEU PRO ALA LEU LEU VAL LEU LEU GLY ALA LEU PHE PRO
 GGA CCT GGC AAT GCC CAG ACA TCT GTG TCC CCC TCA AAA GTC ATC CTG CCC CGG GGA GGC TCC GTG
 GLY PRO GLY ASN ALA GLN THR SER VAL SER PRO SER LYS VAL ILE LEU PRO ARG GLY GLY SER VAL
 CTG GTG ACA TGC AGC ACC TCC TGT GAC CAG CCC AAG TTG TGG GGC ATA GAG ACC CCG TTG CCT AAA
 LEU VAL THR CYS SER THR SER CYS ASP GLN PRO LYS LEU LEU GLY ILE GLU THR PRO LEU PRO LYS
 AAG GAG TTG CTC CTG CCT GGG AAC AAC CGG AAG GTG TAT GAA CTG AGC AAT GTG CAA GAA GAT AGC
 LYS GLU LEU LEU PRO GLY ASN ASN ARG LYS VAL TYR GLU LEU SER ASN VAL GLN GLU ASP SER
 CAA CCA ATG TGC TAT TCA AAC TGC CCT GAT GGG CAG TCA ACA GCT AAA ACC TTC CTC ACC GTG TAC
 GLN PRO MET CYS TYR SER SER ASN CYS PRO ASP GLY GLN SER THR ALA LYS THR PHE LEU THR VAL TYR
 TGG ACT CCA GAA CGG GTG GAA CTG GCA CCC CTC CCC TCT TGG CAG CCA GTG GGC AAG AAC CTT ACC
 TRP THR PRO PRO GLU ARG VAL GLU LEU ALA PRO LEU PRO SER TRP GLN PRO VAL GLY LYS ASN LEU THR
 CTA CGC TGC CAG GTG GAG GGT GGG GCA CCC CGG GCC AAC CTC ACC GTG GTG CTC GTG GGG GAG
 LEU ARG CYS GLN VAL GLU GLY GLY ALA PRO ARG ALA ASN LEU THR VAL VAL LEU ARG GLY GLU
 AAG GAG CTG AAA CGG GAG CCA GCT GTG GGG GAG CCC GCT GAG GTC ACG ACC ACG GTG CTG GTG AGG
 LYS GLU LEU LYS ARG GLU PRO ALA VAL GLY GLU PRO ALA GLU VAL THR THR THR VAL LEU VAL ARG
 AGA GAT CAC CAT GGA GCC AAT TTC TCG TGC CGC ACT GAA CTG GAC CTG CGG CCC CAA GGG CTG GAG
 ARG ASP HIS HIS GLY ALA ASN PHE SER CYS ARG THR GLU LEU ASP LEU ARG PRO GLN GLY LEU GLU

FIG. 1A

2/26

CTG TTT GAG AAC ACC TCG GCC TCG CCC TAC CAG CTC CAG ACC TTT GTC CTG CCA GCG ACT CCC CCA CAA
 LEU PHE GLU ASN THR SER ALA PRO TYR GLN LEU GLN THR PHE VAL GTC CTG TCC CTG GAC GGG CTG
 CTT GTC AGC CCC CGG GTC CTA GAG GTG GAC ACG CAG CAG GGG ACC GTG GTC TGT TCC CTG GAC GGG CTG
 LEU VAL SER PRO ARG VAL LEU GLU VAL ASP THR GLN GLN GLY THR VAL VAL CYS SER LEU ASP GLY LEU
 TTC CCA GTC TCG GAG GCC CAG GTC CAC CTG GCA CTG GCG GAC CAG AGG TTG AAC CCC ACA GTC ACC
 PHE PRO VAL SER GLU ALA GLN VAL HIS LEU ALA LEU GLY ASP GLN ARG LEU ASN PRO THR VAL THR
 TAT GGC AAC GAC TCC TTC TCG GCC AAG GCC TCA GTC AGT GTG ACC GCA GAG GAC GAG GGC ACC CAG
 TYR GLY ASN ASP SER PHE SER ALA LYS ALA SER VAL SER VAL THR ALA GLU ASP GLU GLY THR GLN
 CGG CTG ACG TGT GCA GTA ATA CTG GGG AAC CAG AGC CAG GAG ACA CTG CAG ACA GTG ACC ATC TAC
 ARG LEU THR CYS ALA VAL ILE LEU GLY ASN GLN SER GLN GLU THR LEU GLN THR VAL THR ILE TYR
 AGC TTT CCG GCG CCC AAC GTG ATT CTG ACG AAG CCA GAG GTC TCA GAA GGG ACC GAG GTG ACA GTG
 SER PHE PRO ALA PRO ASN VAL ILE LEU THR LYS PRO GLU VAL SER GLU GLY THR GLU VAL THR VAL
 AAG TGT GAG GCC CAC CCT AGA GCC AAG GTG ACG CTG AAT GGG GTT CCA GCC CAG CCA CTG GGC CCG
 LYS CYS GLU ALA HIS PRO ARG ALA LYS VAL THR LEU ASN GLY ARG SER PHE SER CYS SER ALA THR
 AGG GCC CAG CTC CTG AAG GCC ACC CCA GAG GAC AAC GCG GCG AGC TTC TCC TGC TCT GCA ACC
 ARG ALA GLN LEU LEU LYS ALA THR PRO GLU ASP ASN GLY ARG SER PHE SER CYS SER ALA THR
 CTG GAG GTG GCC GGC CAG CTT ATA CAC AAG AAC CAG ACC CGG GAG CTT CGT GTC CTG TAT GGC CCC
 LEU GLU VAL ALA GLY GLN LEU ILE HIS LYS ASN GLN THR ARG GLU LEU ARG VAL LEU TYR GLY PRO
 CGA CTG GAC GAG AGG GAT TGT CCG GGA AAC TGG ACG TGG CCA GAA AAT TCC CAG CAG ACT CCA ATG
 ARG LEU ASP GLU ARG ASP CYS PRO GLY ASN TRP THR TRP PRO GLU ASN SER GLN GLN THR PRO MET
 TGC CAG GCT TGG GGG AAC CCA TTG CCC GAG CTC AAG TGT CTA AAG GAT GGC ACT TTC CCA CTG CCC
 CYS GLN ALA TRP GLY ASN PRO LEU PRO GLU LEU LYS CYS LEU LYS ASP GLY THR PHE PRO LEU PRO

FIG. 1B

3/26

ATC GGG GAA TCA GTG ACT GTC ACT CGA GAT CTT GAG GGC ACC TAC CTC TGT CGG GCC AGG AGC ACT
 ILE GLY GLU SER VAL THR VAL THR ARG ASP LEU LEU GLU GLY THR TYR LEU CYS ARG ALA ARG SER THR

 CAA GGG GAG GTC ACC CGC GAG GTG ACC GTG AAT GTG CTC TCC CCC CGG TAT GAG ATT GTC ATC ATC
 GLN GLY GLU VAL THR ARG GLU VAL THR VAL ASN VAL LEU SER PRO ARG TYR GLU ILE VAL ILE ILE

 ACT GTG GTA GCA GCC GCA GTC ATA ATG GGC ACT GCA GGC CTC AGC ACG TAC CTC TAT AAC CGC CAG
 THR VAL VAL ALA ALA VAL ILE MET GLY THR ALA GLY LEU SER THR TYR LEU TYR ASN ARG GLN

 CGG AAG ATC AAG AAA TAC AGA CTA CAA CAG GCC CAA AAA GGG ACC CCC ATG AAA CCG AAC ACA CAA
 ARG LYS ILE LYS LYS TYR ARG LEU GLN ALA GLN LYS GLY THR PRO MET LYS PRO ASN THR GLN

 GCC ACG CCT CCC TGA ACCTATCCCG GGACAGGGCC TCTTCTCTCGG CCTTCCCATA TTGGTGGCAG TGGTGCCACA
 ALA THR PRO PRO ***

 CTGAACAGAG TGAAGACAT ATGCCATGCA GCTACACCTA CCGGCCCTGG GACGCCGGAG GACAGGGCAT TGTCTCAGT
 CAGATACAAC AGCATTTGGG GCCATGGTAC CTGCACACCT AAAACACTAG GCCACGCATC TGATCTGTAG TCACATGACT
 AAGCCAAGAG GAAGGAGCAA GACTCAAGAC ATGATTGATG GATGTTAAAG TCTAGCCTGA TGAGAGGGGA AGTGGTGGG
 GAGACATAGC CCCACCATGA GGACATACAA CTGGGAAATA CTGAAACTTG CTGCCTATTG GGTATGCTGA GGCCACACAGA
 CTTACAGAAG AAGTGGCCCT CCATAGACAT GTGTAGCATC AAAACACAAA GGCCACACT TCCTGACGGA TGCCAGCTTG
 GGCACCTGCTG TCTACTGACC CCAACCCCTTG ATGATATGTA TTTATTCAAT TGTATTTTA CCAGCTATT ATTGAGTGTC
 TTTTATGTAG GCTAAATGAA CATAGGTCTC TGGCCTCAGG GAGCTCCCAG TCCATGTAC ATTCAAGGTC ACCAGGTACA
 GTTGACAGG TTGTACACTG CAGGAGAGTG CCTGGCAAAA AGATCAAATG GGGCTGGGAC TTCTCATTTGG CCAACCTGCC
 TTTCCCCAGA AGGAGTGATT TTTCTATCGG CACAAAAGCA CTATATGGAC TGGTAATGTT TCACAGGTTT AGAGATTACC

FIG. 1C

4/26

CAGTGAGGCC TTATTCTCTCC CTTCCCCCCA AAACTGACAC CTTTGTTAGC CACCTCCCCA CCCACATACA TTTCTGCCAG
TGTTACAATG ACACTCAGCG GTCATGCTG GACATGAGTG CCCAGGGAAT ATGCCCCAAGC TATGCCTTGT CCTCTTGTC
TGTTGCAAT TCACTGGGAG CTTGCACTAT TGCAGCTCCA GTTTCCTGCA GTGATCAGGG TCCTGCAAGC AGTGGGGAAG
GGGGCCAAGG TATTGGAGGA CTCCTCCCCA GCTTTGGAAG GGTCAATCCG GTGTGTGTGT GTGTGTATGT GTAGACAAGC
TCTCGCTCTG TCACCCAGGC TGGAGTGCAG TGGTGCAATC ATGTTCACT GCAGTCTTGA CCTTTGGGC TCAAGTGATC
CTCCACCTC AGCCTCCTGA GTAGTGGA CCATAGGCTC ACAACACCCAC ACCTGGCAAA TTTGATTTTT TTTTTTTTT
TCAGAGACGG GGTCTCGCAA CATTGCCCAG ACTTCCTTTG TGTTAGTTAA TAAAGCTTTC TCAACTGCCA AAAAAAAAAA
AAAAAA

FIG. 1D

5/26

FIG. 2A

TTCACATCAA AACTCCTATA CTGACCTGAG ACAGAGGCAG CAGTGATACC CACCTGAGAG ATCCTGTGTT TGA
 ACAACTG CTTCCCAAAA CGGAAAGTAT TTCAAGCCTA AACCTTTGGG TGAAGAAGAC TCTTGAAGTC ATG ATT
 met ile
 GCT TCA CAG TTT CTC TCA GCT CTC ACT TTG GTG CTT CTC ATT AAA GAG AGT GGA GCC TGG
 ala ser gln phe leu ser ala leu thr leu val leu leu ile lys glu ser gly ala trp
 TCT TAC AAC ACC TCC ACG GAA GCT ATG ACT TAT GAT GAG GCC AGT GCT TAT TGT CAG CAA
 ser tyr asn thr ser thr glu ala met thr tyr asp glu ala ser ala tyr cys gln gln
 AGG TAC ACA CAC CTG GTT GCA ATT CAA AAC AAA GAA GAG ATT GAG TAC CTA AAC TCC ATA
 arg tyr thr his leu val ala ile gln asn lys glu glu ile glu tyr leu asn ser ile
 TTG AGC TAT TCA CCA AGT TAT TAC TGG ATT GGA ATC AGA AAA GTC AAC AAT GTG TGG GTC
 leu ser tyr ser pro ser tyr tyr trp ile gly ile arg lys val asn asn val trp val
 TGG GTA GGA ACC CAG AAA CCT CTG ACA GAA GAA GCC AAG AAC TGG GCT CCA GGT GAA CCC
 trp val gly thr gln lys pro leu thr glu glu ala lys asn trp ala pro gly glu pro
 AAC AAT AGG CAA AAA GAT GAG GAC TGC GTG GAG ATC TAC ATC AAG AGA GAA AAA GAT GTG
 asn asn arg gln lys asp glu asp cys val glu ile tyr ile lys arg glu lys asp val
 GCC ATG TGG AAT GAT GAG AGG TGC AGC AAG AAG CTT GCC CTA TGC TAC ACA GCT GCC
 gly met trp asn asp glu arg cys ser lys lys leu ala leu cys tyr thr ala ala
 TGT ACC AAT ACA TCC TGC AGT GGC CAC GGT GAA TGT GTA GAG ACC ATC AAT AAT TAC ACT
 cys thr asn thr ser cys ser gly his gly glu cys val glu thr ile asn asn tyr thr
 TGC AAG TGT GAC CCT GGC TTC AGT GGA CTC AAG TGT GAG CAA ATT GTG AAC TGT ACA GCC
 cys lys cys asp pro gly phe ser gly leu lys cys glu gln ile val asn cys thr ala

6/26

CTG GAA TCC CCT GAG CAT GGA AGC CTG GTT TGC AGT CAC CCA CTG GGA AAC TTC AGC TAC
 leu glu ser pro glu his gly ser leu val cys ser his pro leu gly asn phe ser tyr

AAT TCT TCC TGC TCT ATC AGC TGT GAT AGG GGT TAC CTG CCA AGC AGC ATG GAG ACC ATG
 asn ser ser cys ser ile ser cys asp arg gly tyr leu pro ser ser met glu thr met

CAG TGT ATG TCC TCT GGA GAA TGG AGT GCT CCT ATT CCA GCC TGC AAT GTG GTT GAG TGT
 gln cys met ser ser gly glu trp ser ala pro ile pro ala cys asn val val glu cys

GAT GCT GTG ACA AAT CCA GCC AAT GGG TTC GTG GAA TGT TTC CAA AAC CCT GGA AGC TTC
 asp ala val thr asn pro ala asn gly phe val glu cys phe gln asn pro gly ser phe

CCA TGG AAC ACA ACC TGT ACA TTT GAC TGT GAA GAA GGA TTT GAA CTA ATG GGA GCC CAG
 pro trp asn thr thr cys thr phe asp cys glu glu gly phe glu leu met gly ala gln

AGC CTT CAG TGT ACC TCA TCT GGG AAT TGG GAC AAC GAG AAG CCA ACG TGT AAA GCT GTG
 ser leu gln cys thr ser ser gly asn trp asp asp asn glu lys pro thr cys lys ala val

ACA TGC AGG GCC GTC CGC CAG CCT CAG AAT GGC TCT GTG AGG TGC AGC CAT TCC CCT GCT
 thr cys arg ala val arg gln pro gln asn gly ser val arg cys ser his ser pro ala

GGA GAG TTC ACC TTC AAA TCA TCC TGC AAC TTC ACC TGT GAG GAA GGC TTC ATG TTG CAG
 gly glu phe thr phe lys ser ser cys asn phe thr cys glu glu gly phe met leu gln

GGA CCA GCC CAG GTT GAA TGC ACC ACT CAA GGG CAG TGG ACA CAG CAA ATC CCA GTT TGT
 gly pro ala gln val glu cys thr thr gln gly gln trp thr gln gln ile pro val cys

GAA GCT TTC CAG TGC ACA GCC TTG TCC AAC CCC GAG CGA GGC TAC ATG AAT TGT CTT CCT
 glu ala phe gln cys thr ala leu ser asn pro glu arg gly tyr met asn cys leu pro

FIG. 2B

7/26

AGT GCT TCT GGC AGT TTC CGT TAT GGG TCC AGC TGT GAG TTC TCC TGT GAG CAG GGT TTT
 ser ala ser gly ser phe arg tyr gly ser ser cys cys glu phe ser cys gly phe
 GTG TTG AAG GGA TCC AAA AGG CTC CAA TGT GGC CCC ACA GGG GAG TGG GAC AAC GAG AAG
 val leu lys gly ser lys arg leu gln cys gly pro thr gly glu trp asp asn glu lys
 CCC ACA TGT GAA GCT GTG AGA TGC GAT GCT GTC CAC CAG CCC CCG AAG GGT TTG GTG AGG
 pro thr cys glu ala val arg cys asp ala val his gln pro pro lys gly leu val arg
 TGT GCT CAT TCC CCT ATT GGA GAA TTC ACC TAC AAG TCC TCT TGT GCC TTC AGC TGT GAG
 cys ala his ser pro ile gly glu phe thr tyr lys ser ser cys ala phe ser cys glu
 GAG GGA TTT GAA TTA TAT GGA TCA ACT CAA CTT GAG TGC ACA TCT CAG GGA CAA TGG ACA
 glu gly phe glu leu tyr gly ser thr gln leu glu cys thr ser gln gly gln trp thr
 GAA GAG GTT CCT TCC TGC CAA GTG GTA AAA TGT TCA AGC CTG GCA GTT CCG GGA AAG ATC
 glu glu val pro ser cys gln val val lys cys ser ser leu ala val pro gly lys ile
 AAC ATG AGC TGC AGT GGG GAG CCC GTG TTT GGC ACT GTG TGC AAG TTC GCC TGT CCT GAA
 asn met ser cys ser gly glu pro val phe gly thr val cys lys phe ala cys pro glu
 GGA TGG ACG CTC AAT GGC TCT GCA GCT CGG ACA TGT GGA GCC ACA GGA CAC TGG TCT GGC
 gly trp thr leu asn gly ser ala ala arg thr cys gly ala thr gly his trp ser gly
 CTG CTA CCT ACC TGT GAA GCT CCC ACT GAG TCC AAC ATT CCC TTTG GTA GCT GGA CTT TCT
 leu leu pro thr cys glu ala pro thr glu ser asn ile pro leu val ala gly leu ser
 GCT GCT GGA CTC TCC CTC CTG ACA TTA GCA CCA TTT CTC CTC TGG CTT CGG AAA TGC TTA
 ala ala gly leu ser leu thr leu ala pro phe leu leu trp leu arg lys cys leu
 CGG AAA GCA AAG AAA TTT GTT CCT GCC AGC TGC CAA AGC CTT GAA TCA GAC GGA AGC
 arg lys ala lys lys phe val pro ala ser ser cys gln ser leu glu ser asp gly ser

FIG. 2C

8/26

TAC CAA AAG CCT TCT TAC ATC CTT TAA GTTCAAA AGAATCAGAA ACAGGTGCAT CTGGGGAACT A
tyr gln lys pro ser tyr ile leu ***

GAGGGATAC ACTGAAGTTA ACAGAGACAG ATAACCTCTCC TCGGGTCTCT GGCCCTTCTT GCCTACTATG CCAG
ATGCCCT TTATGGCTGA AACCGCAACA CCCATCACCA CTTCAATAGA TCAAAGTCCA GCAGGCAAGG ACGGCCCT
TCA ACTGAAAAGA CTCAGTGTTT CCTTTCCTAC TCTCAGGATC AAGAAAGTGT TGGCTAATGA AGGAAAGGA
TATTTTCTTC CAAGCAAAGG TGAAGAGACC AAGACTCTGA AATCTCAGAA TTCCTTTTCT AACTCTCCCT TG
CTCGCTGT AAAATCTTGG CACAGAAACA CAATATTTTG TGGCTTCTCT TCTTTTGCCC TTCACAGTGT TTCGA
CAGCT GATTACACAG TTGCTGTCTAT AAGAATGAAT AATAATTATC CAGAGTTTAG AGGAAAAAAA TGAATAAA
AA TATTATAACT TAAAAAATG ACAGATGTTG AATGCCCCACA GGCAAAATGCA TGGAGGGTTG TTAATGGTGC
AAATCCTACT GAATGCTCTG TCGAGGGGTT ACTATGCACA ATTTAATCAC TTTCATCCCT ATGGGATTCA GTG
CTTCTTA AAGAGTTCTT AAGGATGTG ATATTTTAC TTGCATTGAA TATATTATAA TCTTCCATAC TTCCTTC
ATTC AATACAAGTG TGGTAGGGAC TTAAAAAACT TGTAATGCT GTCAACTATG ATATGGTAA AGTTACTTA
T TCTAGATTAC CCCCTCATG TTTATTAACA AATTATGTTA CATCTGTTTT AAATTATTTT CAAAAAGGGA A
ACTATTGTC CCCTAGCAAG GCATGATGTT AACCAGAATA AAGTTCTGAG TGTTTTACT ACAGTTGTTT TTTG
AAAACA TGGTAGAATT GGAGAGTAAA AACTGAATGG AAGGTTTGT TATTGTCAGA TATTTTTC GAAATAT
GTG GTTTCACCA TGA AAAA ACTT CCATGAGGCC AAACGTTTG AACTAATAAA AGCATAAATG CAAACACACA
AAGGTATAAT TTTATGAATG TCTTTGTGG AAAAGAATAC AGAAGATGG ATGTGCTTG CATTCCTACA AA
GATGTTTG TCAGATGTGA TATGTAAACA TAATTCTTGT ATATTATGGA AGATTTTAA TTCACAATAG AAAC

FIG. 2D

SUBSTITUTE SHEET

9/26

CACCA TGTAAAGAG TCATCTGGTA GATTTTAAAC GAATGAAGAT GTCTAATAGT TATTCCTAT TTGTTTTC
TT CTGTATGTTA GGGTGCTCTG GAAGAGAGGA ATGCCTGTGT GAGCAAGCAT TTATGTTTAT TTATAAGCAG
ATTTAACAAAT TCCAAAGGAA TCTCCAGTTT TCAGTTGATC ACTGGCAATG AAAAATTCTC AGTCAGTAAT TGC
CAAAGCT GCTCTAGCCT TGAGGAGTGT GAGAAATCAAA ACTCTCCTAC ACTTCCATTA ACTTAGCATG TGTGTA
AAAA AAAAGTTTCA GAGAAATTCT GGCTGAACAC TGGCAACGAC AAAGCCCAACA GTCAAAACAG AGATGTGAT
A AGGATCAGAA CAGCAGAGGT TCTTTTAAAG GGCAGAAAA ACTCTGGGA ATAAGAGAGA ACAACTACTG T
GATCAGGCT ATGTATGGAA TACAGTGTTA TTTTCTTTGA AATTGTTTAA GTGTTGTAAA TATTTATGTA AACT
GCATTA GAAATTAGCT GTGTGAAATA CCAGTGTGGT TTGTGTTTGA GTTTTATTGA GAATTTTAA TTATAAC
TTA AAATATTTTA TAATTTTAA AGTATATATT TATTTAAGCT TATGTCAGAC CTATTTGACA TAACACTATA
AAGGTTGACA ATAAATGTGC TTATGTTT

FIG. 2E

10/26

FIG. 3A

CGGGCCTCAC TGGCTTCAGG AGCTGAATAC CCTCCAGGC ACACACAGGT GGGACACAAA TAAGGTTT GGA
 ACCACTA TTTTCTCATC ACGACAGCAA CTTAAA ATG CCT GGG AAG ATG GTC GTG ATC CTT GGA GCC
 met pro gly lys met val val ile leu gly ala
 TCA AAT ATA CTT TGG ATA ATG TTT GCA GCT TCT CAA GCT TTT AAA ATC GAG ACC ACC CCA
 ser asn ile leu trp ile met phe ala ala ser gln ala phe lys ile glu thr thr pro
 GAA TCT AGA TAT CTT GCT CAG ATT GGT GAC TCC GTC TCA TTG ACT TGC AGC ACC ACA GGC
 glu ser arg tyr leu ala gln ile gly asp ser val ser leu thr cys ser thr thr gly
 TGT GAG TCC CCA TTT TTC TCT TGG AGA ACC CAG ATA GAT AGT CCA CTG AAT GGG AAG GTG
 cys glu ser pro phe ser trp arg thr gln ile asp ser pro leu asn gly lys val
 ACG AAT GAG GGG ACC ACA TCT ACG CTG ACA ATG AAT CCT GTT GGT GGG AAC GAA CAC
 thr asn glu gly thr ser thr ser leu thr met asn pro val ser phe gly asn glu his
 TCT TAC CTG TGC ACA GCA ACT TGT GAA TCT AGG AAA TTG GAA AAA GGA ATC CAG GTG GAG
 ser tyr leu cys thr ala thr cys glu ser arg lys leu glu lys gly ile gln val glu
 ATC TAC TCT TTT CCT AAG GAT CCA GAG ATT CAT TTG AGT GGC CCT CTG GAG GCT GGG AAG
 ile tyr ser phe pro lys asp pro glu ile his leu ser gly pro leu glu ala gly lys
 CCG ATC ACA GTC AAG TGT TCA GTT GCT GAT GTA TAC CCA TTT GAC AGG CTG GAG ATA GAC
 pro ile thr val lys cys ser val ala asp val tyr pro phe asp arg leu glu ile asp
 TTA CTG AAA GGA GAT CAT CTC ATG AAG AGT CAG GAA TTT CTG GAG GAT GCA GAC AGG AAG
 leu leu lys gly asp his leu met lys ser gln glu phe leu glu asp ala asp arg lys
 TCC CTG GAA ACC AAG AGT TTG GAA GTA ACC TTT ACT CCT GTC ATT GAG GAT ATT GGA AAA
 ser leu glu thr lys ser leu glu val thr phe thr pro val ile glu asp ile gly lys
 GTT CTT GTT TGC CGA GCT AAA TTA CAC ATT GAT GAA ATG GAT TCT GTG CCC ACA GTA AGG
 val leu val cys arg ala lys leu his ile asp glu met asp ser val pro thr val arg

11/26

CAG GCT GTA AAA GAA TTT CAA GTC TAC ATA TCA CCC AAG AAT ACA GTT ATT TCT GTG AAT
 gln ala val lys glu leu gln val tyr ile ser pro lys asn thr val ile ser val asn

 CCA TCC ACA AAG CTG CAA GAA GGT GGC TCT GTG ACC ATG ACC TGT TCC AGC GAG GGT CTA
 pro ser thr lys leu gln glu gly gly ser val thr met thr cys ser ser glu gly leu

 CCA GCT CCA GAG ATT TTC TGG AGT AAG AAA TTA GAT AAT GGG AAT CTA CAG CAC CTT TCT
 pro ala pro glu ile phe trp ser lys lys leu asp asn gly asn leu gln his leu ser

 GGA AAT GCA ACT CTC ACC TTA ATT GCT ATG AGG ATG GAA GAT TCT GGA ATT TAT GTG TGT
 gly asn ala thr leu thr leu ile ala met arg met glu asp ser gly ile tyr val cys

 GAA GGA GTT AAT TTG ATT GGG AAA AAC AGA AAA GAG GTG GAA TTA ATT GTT CAA GCA TTC
 glu gly val asn leu ile gly lys asn arg lys glu val glu leu ile val gln ala phe

 CCT AGA GAT CCA GAA ATC GAG ATG AGT GGT GGC CTC GTG AAT GGG AGC TCT GTG ACT GTA
 pro arg asp pro glu ile glu met ser gly gly leu val asn gly ser val thr val

 AGC TGC AAG GTT CCT AGC GTG TAC CCC CTT GAC CCG CTG GAG ATT GAA TTA CTT AAG GGG
 ser cys lys val pro ser val tyr pro leu asp arg leu glu ile glu leu lys gly

 GAG ACT ATT CTG GAG AAT ATA GAG TTT TTG GAG GAT ACG GAT ATG AAA TCT CTA GAG AAC
 glu thr ile leu glu asn ile glu phe leu glu asp thr asp met lys ser leu glu asn

 AAA AGT TTG GAA ATG ACC TTC ATC CCT ACC ATT GAA GAT ACT GGA AAA GCT CTT GTT TGT
 lys ser leu glu met thr phe ile pro thr ile glu asp thr gly lys ala leu val cys

 CAG GCT AAG TTA CAT ATT GAT GAC ATG GAA TTC GAA CCC AAA CAA AGG CAG AGT ACG CAA
 gln ala lys leu his ile asp asp met glu phe glu pro lys gln arg gln ser thr gln

 ACA CTT TAT GTC AAT GTT GCC CCC AGA GAT ACA ACC GTC TTG GTC AGC CCT TCC TCC ATC
 thr leu tyr val asn val ala pro arg asp thr thr val leu val ser pro ser ser ile

 CTG GAG GAA GGC AGT TCT GTG AAT ATG ACA TGC TTG AGC CAG GGC TTT CCT GCT CCG AAA
 leu glu glu gly ser ser val asn met thr cys leu ser gln gly phe pro ala pro lys

FIG. 3B

12/26

ATC CTG TGG AGC AGG CAG CTC CCT AAC GGG GAG CTA CAG CCT CTT TCT GAG AAT GCA ACT
 ile leu trp ser arg gln leu pro asn gly glu leu gln pro leu ser glu asn ala thr
 CTC ACC TTA ATT TCT ACA AAA ATG GAA GAT TCT GGG GTT TAT TTA TGT GAA GGA ATT AAC
 leu thr leu ile ser thr lys met glu asp ser gly val tyr leu cys glu gly ile asn
 CAG GCT GGA AGA AGC AGA AAG GAA GTG GAA TTA ATT ATC CAA GTT ACT CCA AAA GAC ATA
 gln ala gly arg ser arg lys glu val glu leu ile ile gln val thr pro lys asp ile
 AAA CTT ACA GCT TTT CCT TCT GAG AGT GTC AAA GAA GGA GAC ACT GTC ATC ATC TCT TGT
 lys leu thr ala phe pro ser glu ser val lys glu gly asp thr val ile ile ser cys
 ACA TGT GGA AAT GTT CCA GAA ACA TGG ATA ATC CTG AAG AAA AAA GCG GAG ACA GGA GAC
 thr cys gly asn val pro glu thr trp ile ile leu lys lys ala glu thr gly asp
 ACA GTA CTA AAA TCT ATA GAT GGC GCC TAT ACC ATC CGA AAG GCC CAG TTG AAG GAT GCG
 thr val leu lys ser ile asp gly ala tyr thr ile arg lys ala gln leu lys asp ala
 GGA GTA TAT GAA TGT GAA TCT AAA AAC AAA GTT GGC TCA CAA TTA AGA AGT TTA ACA CTT
 gly val tyr glu cys glu ser lys asn lys val gly ser gln leu arg ser leu thr leu
 GAT GTT CAA GGA AGA GAA AAC AAC AAA GAC TAT TTT TCT CCT GAG CTT CTC GTG CTC TAT
 asp val gln gly arg glu asn asn lys asp tyr phe ser pro glu leu val leu tyr
 TTT GCA TCC TCC TTA ATA ATA CCT GCC ATT GGA ATG ATA ATT TAC TTT GCA AGA AAA GCC
 phe ala ser ser leu ile ile pro ala ile gly met ile ile tyr phe ala arg lys ala
 AAC ATG AAG GGG TCA TAT AGT CTT GTA GAA GCA CAG AAA TCA AAA GTG TAG CTAATGCTTG
 asn met lys gly ser tyr ser leu val glu ala gln lys ser lys val ***
 ATATGTTCAA CTGGAGACAC TATTTATCTG TGCAAATCCT TGATACTGCT CATCATTCCT TGAGAAAAAC AAT
 GAGCTGA GAGGCAGACT TCCCTGAATG TATTGAACCTT GGAAGAANAAT GCCCATCTAT GTCCCTTGCT GTGAGC
 AAGA AGTCAAAGTA AAACCTTGCTG CCTGAAGAAC AGTAACTGCC ATCAAGATGA GAGAACTGGA GGAGTTCCT
 T GATCTGTATA TACAATAACA TAATTGTGAC ATATGTAAAA TAAATATTG CCATAGCAAG ATTGCTTAAAA

FIG. 3C

13/26

TAGCAACAC TCTATATTTA GATTGTTAAA ATAACTAGTG TTGCTTGGAC TATTATAATT TAATGCATGT TAGG
AAAATT TCACATTAAAT ATTTGCTGAC AGCTGACCCTT TGTCACTCTTT CTTCATATTTT ATTCCCTTTC ACAAAAT
TTT ATTCCTATAT AGTTTATTGA CAATAATTTC AGGTTTTGTA AAGATGCCCGG GTTTTATATT TTTATAGACA
AATAATAAGC AAAGGGAGCA CTGGGTTGAC TTTCAGGTAC TAAATACCTC AACCTATGGT ATAATGGTTG AC
TGGGTTTC TCTGTATAGT ACTGGCATGG TACGGAGATG TTTCACGAAG TTTGTTCATC AGACTCCCTGT GCAAC
TTTCC CAATGTGGCC TAAAAATGCA ACTTCCTTTT ATTTCTCTTT GTAAATGTTT AGGTTTTTTT GTATAGTA
AA GTGATAATTT CTGGAATTAA AAA

FIG. 3D

14/26




FIG. 4

15/26




FIG. 5

16/26




FIG. 6A




FIG. 6B

17/26




FIG. 7

18/26




FIG. 8

19/26




FIG. 9

20/26




FIG. 10

21/26





FIG. 11

22/26



• 2679 + 2674 * 2673 □ 2687
× 2686 ◇ 2683 △ 1571 (ICAM-1)

FIG. 12




FIG. 13

24/26




FIG. 14

25/26





FIG. 15

26/26

**FIG. 16**

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US93/08101

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A61K 48/00

US CL : 514/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Dialog & APS "hybridization", "triple-stranded", ICAMM-1 "ECAM-1", "VCAM-1", "antisense", therapy

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|---------------|--|---|
| <u>X</u> Y | Chemical Reviews, volume 90, no. 4, issued June 1990, Uhlmann <u>et al.</u> , "Antisense Oligonucleotides: A new therapeutic principle", pages 544-584, see entire article. | <u>1-5, 9-14</u> 16-18, 26-30, 34-36, 38, 39, 53 |
| <u>X</u> Y | Journal of Biological Chemistry, volume 266, no. 27, issued 25 September 1991, Chiang <u>et al.</u> , "Antisense oligonucleotides inhibit intercellular adhesion molecule 1 expression by two distinct mechanisms", pages 18162-18171, see entire article. | 1, 2, 4, 6, 15, 19- <u>22, 25, 45, 46, 50</u> 7, 8, 31-33, 37, 44, 56-59, 64-66, 69-72, 74, 76 |

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

| | | |
|---|-----|--|
| * Special categories of cited documents: | *T | later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be part of particular relevance | *X* | document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *E* earlier document published on or after the international filing date | *Y* | document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *A* | document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | | |
| *P* document published prior to the international filing date but later than the priority date claimed | | |

Date of the actual completion of the international search

16 NOVEMBER 1993

Date of mailing of the international search report

DEC 02 1993

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

SUZANNE ZISKA

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/08101

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|---|--|
| Y | Cell, Volume 59, issued 22 December 1989, Osborn <u>et al.</u> , "Direct expression cloning of vascular cell adhesion molecule 1, a cytokine-induced endothelial protein that binds to lymphocytes," pages 1203-1211, see entire article. | 24, 42, 43, 49, 52, 55, 68 |
| Y | Science, volume 246, issued 08 December 1989, Rice <u>et al.</u> , "An inducible endothelial cell surface glycoprotein mediates melanoma adhesion", pages 1303-1306, see entire article. | 23, 40, 41, 47, 48, 51, 54, 60- 63, 67, 73, 75 |
| Y | Science, volume 243, issued 03 March 1989, Bevilacqua <u>et al.</u> , "Endothelial leukocyte adhesion molecule 1: AN inducible receptor for neutrophils related to complement regulatory proteins and lectins", pages 1160-1165, see entire article. | 23, 40, 41, 47, 48, 51, 54, 60- 63, 67, 73, 75 |

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows: 1

I. Claims 1-27, 40-43 and 45-52, drawn to a first product, an oligonucleotide, classified in Class 536, subclass 24.5, for example, and a first method of using the first product, classified in 435, subclass 172.3, for example;

II. Claims 28-39, 44, 53-62, drawn to a second method of using the first product, a method of treating an animal comprising contacting an animal with a therapeutically effective amount of an oligonucleotide, classified in Class 514, subclass 44, for example;

III. Claims 63-73, drawn to a third method of using the first product, a method of decreasing metastasis in an animal comprising contacting an animal with a therapeutically effective amount of an oligonucleotide, classified in Class 514, subclass 44, for example;

IV. Claims 74-76, drawn to a fourth method of using the first product, a method of treating psoriasis comprising contacting an animal with a therapeutically effective amount of an oligonucleotide, classified in Class 514, subclass 44, for example.

The inventions listed as Groups I-IV do not meet the requirements for Unity of Invention for the following reasons:

Each grouping of claims forms a separate invention not linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept. PCT Rules 13.1 and 13.2 do not permit multiple distinct products and methods within a single inventive concept.